

Existence of Solutions of a (Nonsmooth) Thermoviscoplastic Model and Associated Optimal Control Problems

Roland Herzog^a, Christian Meyer^b, Ailyn Schäfer^a

^a*Technische Universität Chemnitz, Faculty of Mathematics, Professorship Numerical
Mathematics (Partial Differential Equations), D-09107 Chemnitz, Germany*

^b*Technische Universität Dortmund, Faculty of Mathematics, Lehrstuhl LSX, D-44227
Dortmund, Germany*

Abstract

A thermoviscoplastic model with linear kinematic hardening, von Mises yield condition and mixed boundary conditions is considered. The existence of a unique weak solution is proved by means of a fixed-point argument, and by employing maximal parabolic regularity theory. The weak continuity of the solution operator is also shown. As an application, the existence of a global minimizer of a class of optimal control problems is proved.

Keywords: thermoviscoplasticity; variational inequality of second kind; mixed boundary conditions; Banach fixed-point theorem; maximal parabolic regularity; optimal control

1. Introduction

We consider the following thermovisco(elasto)plastic model with linear kinematic hardening and von Mises yield condition:

$$\text{stress-strain relation: } \boldsymbol{\sigma} = \mathbb{C}(\boldsymbol{\varepsilon}(\mathbf{u}) - \mathbf{p} - \mathbf{t}(\theta)), \quad (1)$$

$$\text{conjugate forces: } \boldsymbol{\chi} = -\mathbb{H} \mathbf{p}, \quad (2)$$

$$\text{viscoplastic flow rule: } \epsilon \dot{\mathbf{p}} + \partial D(\dot{\mathbf{p}}, \theta) \ni [\boldsymbol{\sigma} + \boldsymbol{\chi}], \quad (3)$$

$$\text{balance of momentum: } -\operatorname{div}(\boldsymbol{\sigma} + \gamma \boldsymbol{\varepsilon}(\dot{\mathbf{u}})) = \boldsymbol{\ell}, \quad (4)$$

$$\text{heat equation: } \rho c_p \dot{\theta} - \operatorname{div}(\kappa \nabla \theta) = r + \gamma \boldsymbol{\varepsilon}(\dot{\mathbf{u}}) : \boldsymbol{\varepsilon}(\dot{\mathbf{u}}) + (\boldsymbol{\sigma} + \boldsymbol{\chi}) : \dot{\mathbf{p}} - \theta \mathbf{t}'(\theta) : \mathbb{C}(\boldsymbol{\varepsilon}(\dot{\mathbf{u}}) - \dot{\mathbf{p}}). \quad (5)$$

The unknowns are the stress $\boldsymbol{\sigma}$, back-stress $\boldsymbol{\chi}$, plastic strain \mathbf{p} , displacement \mathbf{u} and temperature θ . Further, \mathbb{C} and \mathbb{H} denote the elastic and hardening

Email addresses: roland.herzog@mathematik.tu-chemnitz.de (Roland Herzog),
cmeyer@math.tu-dortmund.de (Christian Meyer),
ailyn.schaefer@mathematik.tu-chemnitz.de (Ailyn Schäfer)

moduli, respectively. $\varepsilon(\mathbf{u})$ denotes the symmetrized gradient or linearized strain associated with \mathbf{u} . The temperature dependent term $\mathbf{t}(\theta)$ expresses thermally induced stresses. D denotes the dissipation function. The right hand sides ℓ and r represent mechanical and thermal volume and boundary loads, respectively. ϱ , c_p and κ describe the density, specific heat capacity and thermal conductivity of the material. The positive parameters ϵ and γ represent viscous effects in the evolution of the plastic strain and in the balance of momentum. For the derivation of the system (1)–(5) and more on its physical background, we refer to (Ottosen & Ristinmaa, 2005, Chapter 22 and 23).

The analysis of thermoplastic models poses numerous mathematical challenges, mainly due to the low integrability of the nonlinear terms on the right hand side of the heat equation. Several approaches have been considered in the literature to deduce the existence and uniqueness of a solution, and we mention the following.

- [Chelmiński & Racke \(2006\)](#): In this model without viscosity terms the dissipation function is only allowed to depend linearly on the temperature and a simplified mechanical heat source is used which does not account for plastic dissipation and is cut off at large temperatures. The authors use a Yosida regularization to prove the existence of a solution.
- [Bartels & Roubíček \(2008\)](#): The model does not account for hardening and thermal strains, it contains a hyperbolic viscous balance of momentum and a simplified right hand side of the heat equation. The authors prove the existence of a solution in a weak sense via a discretization strategy.
- [Bartels & Roubíček \(2011\)](#): In contrast to [Bartels & Roubíček \(2008\)](#) the authors take into account thermal strains, linear kinematic and isotropic hardening and the same right hand side of the heat equation as in (5) but they consider a temperature independent flow rule. The authors require a growth condition for the heat capacity w.r.t. the temperature to obtain the existence of a solution in a weak sense, again via a discretization procedure.
- [Paoli & Petrov \(2012\)](#): In contrast to our model the authors assume a C^2 regular boundary in addition to homogeneous boundary conditions for the displacement, which leads to better regularity. Moreover, the dissipation function is assumed to be independent of the temperature. The authors use a growth condition for the heat capacity w.r.t. the temperature to show the existence of a solution in a classical sense by means of Schauder’s fixed point theorem.

Our approach is closest to the one in [Paoli & Petrov \(2012\)](#). We emphasize that we admit more general domains and boundary conditions. The overall strategy to show the existence and uniqueness of a solution is an application of Banach’s fixed point theorem, applied to a reduced problem formulated in the temperature variable alone. In order to apply the fixed-point argument, we make use of the theory of maximal parabolic regularity. The same strategy was used

in Hömberg et al. (2009/10) for the analysis and optimal control of a thermistor problem. Furthermore, we focus our discussion on the case of constant heat capacities. We mention that this case is not included in Paoli & Petrov (2012) since a linear growth of the heat capacity is assumed there. In contrast to the linear dependence of the thermal strain on the temperature in Paoli & Petrov (2012), we allow more general thermal strains \mathbf{t} and only assume them to be globally bounded w.r.t. the temperature. This can be achieved w.l.o.g. by a cut-off outside the relevant temperature regime.

Under the assumptions made precise in section 2, our main result is as follows. (We refer the reader to Theorem 8 for a re-iteration of the theorem.)

Theorem (Main Theorem). *There exists $\bar{p} > 2$ such that for all $2 < p \leq \bar{p}$, there exists $\bar{q} > 2$ (depending on p) such that for all $\bar{q} \leq q < \infty$ and sufficiently smooth right hand sides (ℓ, r) and initial conditions $(\mathbf{u}_0, \mathbf{p}_0, \theta_0)$, there exists a unique weak solution $(\mathbf{u}, \mathbf{p}, \theta)$ of (1)–(5) such that*

$$\begin{aligned} \mathbf{u} &\in W^{1,q}(0, T; \mathbf{W}_D^{1,p}(\Omega)), & \mathbf{p} &\in W^{1,q}(0, T; \mathbf{L}^p(\Omega; \mathbb{R}_{sym}^{3 \times 3})) \text{ trace-free,} \\ \theta &\in W^{1, \frac{q}{2}}(0, T; W_\diamond^{-1, v(p)}(\Omega)) \cap L^{\frac{q}{2}}(0, T; W^{1, v(p)}(\Omega)) \end{aligned}$$

with $v(p)$ given in (6). The stress components $\boldsymbol{\sigma}$ and $\boldsymbol{\chi}$ are obtained from (1)–(2).

Note that compared to Bartels & Roubíček (2008, 2011), we obtain solutions of higher regularity, working with a different notion of a solution. Although we hope that the main theorem is a result of independent interest, we consider in this paper also the existence of global minimizers of certain optimal control problems involving (1)–(5). To this end, we prove a result about the weak sequential continuity of the solution map w.r.t. the right hand side data, see Proposition 29. It will be proved using a technique developed in Bartels & Roubíček (2008).

The paper is organized as follows. After an introduction of the exact setting and the detailed assumptions in section 2, the existence and uniqueness of a solution to the system (1)–(5) is shown in section 3. A short roadmap is presented at the beginning of section 3, describing the break up of the proof into smaller parts. In particular, we prove in subsection 3.1 the existence and uniqueness of a solution to system (1)–(4) for a given temperature field, and in subsection 3.2 the contractivity of the fixed-point map on small time intervals is shown, along with a continuation argument. The proof of the main theorem is given in subsection 3.3. Finally, we prove Proposition 29 on the weak sequential continuity of the solution map in section 4, and deduce the existence of a global minimizer of associated optimal control problems as an application.

2. Notation and Assumptions

In the following, Ω denotes a bounded domain in \mathbb{R}^3 and $T > 0$ is a fixed end time point. The spaces $L^p(\Omega)$ and $W^{k,p}(\Omega)$ denote Lebesgue and Sobolev

spaces, respectively. For a Banach space X and its dual space X' , we denote the duality product as $\langle \cdot, \cdot \rangle_X$ or simply $\langle \cdot, \cdot \rangle$ if no ambiguity arises. The norm of X is always denoted as $\|\cdot\|_X$. In the case $X = W^{1,p}(\Omega)$ we denote the dual by $W_\diamond^{-1,p'}(\Omega)$ where $1/p + 1/p' = 1$.

The space $\text{Lin}(X)$ denotes the space of bounded linear functions from X into itself. Furthermore the space $L^p(0, T; X)$ denotes a Bochner space and the space $W^{1,p}(0, T; X)$ is the subset of $L^p(0, T; X)$ such that distributional time derivative of the elements are again in $L^p(0, T; X)$, see, e.g., (Showalter, 1997, Chapter III). The space $W_0^{1,p}(0, T; X)$ denotes the subspace of functions which vanish in $t = 0$.

Vector-valued functions, and spaces containing such functions are written in bold-face notation. The spaces $\mathbb{R}^{3 \times 3}$ and $\mathbb{R}_{\text{sym}}^{3 \times 3}$ represents the (symmetric) 3×3 matrices. Furthermore, $\mathbb{R}_{\text{trace}}^{3 \times 3}$ denotes the symmetric and trace-free 3×3 matrices. For $\mathbf{p}, \mathbf{q} \in \mathbb{R}^{3 \times 3}$, the inner product and the associated Frobenius norm are denoted by $\mathbf{p} : \mathbf{q} := \text{trace}(\mathbf{p}^\top \mathbf{q})$ and $|\mathbf{p}|$, respectively. The symmetrized gradient $\boldsymbol{\varepsilon}(\mathbf{u})$ is defined as $\frac{1}{2}(\nabla \mathbf{u} + \nabla \mathbf{u}^\top)$. The distributional time derivative of a function f defined on $\Omega \times (0, T)$ is denoted by \dot{f} . Further, we denote by g' the Fréchet derivative of a function g defined on \mathbb{R} . Finally, C denotes a generic nonnegative constant and it is written as $C(\cdot)$ to indicate dependencies.

Now we are able to state our assumptions on the quantities in the thermo-viscoplastic model (1)–(5). We begin with the physical constants and functions. We then proceed to make precise the assumptions on the initial conditions and mechanical and thermal loads, and give the weak formulation of the model. We conclude the section with the assumptions on the domain Ω .

Assumption 1.

1. The moduli $\mathbb{C}, \mathbb{H} : \Omega \rightarrow \text{Lin}(\mathbb{R}_{\text{sym}}^{3 \times 3})$ are
 - (a) elements of $L^\infty(\Omega, \text{Lin}(\mathbb{R}_{\text{sym}}^{3 \times 3}))$,
 - (b) symmetric in the sense that

$$\mathbb{C}_{ijkl} = \mathbb{C}_{jikl} = \mathbb{C}_{klij} \quad \text{and} \quad \mathbb{H}_{ijkl} = \mathbb{H}_{jikl} = \mathbb{H}_{klij},$$

- (c) coercive on $\mathbb{R}_{\text{sym}}^{3 \times 3}$ with coercivity constants $\underline{c}, \underline{h} > 0$, i.e.

$$\boldsymbol{\varepsilon} : \mathbb{C}(\mathbf{x}) \boldsymbol{\varepsilon} \geq \underline{c} |\boldsymbol{\varepsilon}|^2 \quad \text{and} \quad \mathbf{p} : \mathbb{H}(\mathbf{x}) \mathbf{p} \geq \underline{h} |\mathbf{p}|^2$$

for all $\boldsymbol{\varepsilon}, \mathbf{p} \in \mathbb{R}_{\text{sym}}^{3 \times 3}$ and almost all $\mathbf{x} \in \Omega$.

2. The temperature dependent initial uni-axial yield stress $\sigma_0 : \mathbb{R} \rightarrow \mathbb{R}$ is
 - (a) positive and belongs to $L^\infty(\mathbb{R})$,
 - (b) Lipschitz continuous, i.e., there exists $L_{\sigma_0} \geq 0$ such that

$$|\sigma_0(\theta_1) - \sigma_0(\theta_2)| \leq L_{\sigma_0} |\theta_1 - \theta_2| \quad \text{for all } \theta_1, \theta_2 \in \mathbb{R}.$$

3. The temperature dependent dissipation function $D : \mathbb{R}_{\text{sym}}^{3 \times 3} \times \mathbb{R} \rightarrow \mathbb{R}$ is defined as

$$D(\mathbf{q}, \theta) := \sqrt{\frac{2}{3}} \sigma_0(\theta) |\mathbf{q}|.$$

4. The temperature dependent thermal strain function $\mathbf{t} : \mathbb{R} \rightarrow \mathbb{R}_{sym}^{3 \times 3}$ is
 - (a) of class $C_b^2(\mathbb{R}, \mathbb{R}_{sym}^{3 \times 3})$ (the space of bounded C^2 functions with bounded derivatives),
 - (b) such that $\mathbb{R} \ni \theta \mapsto \theta \mathbf{t}'(\theta) \in \mathbb{R}_{sym}^{3 \times 3}$ is Lipschitz continuous.
5. The density ϱ , specific heat capacity c_p , thermal conductivity κ and heat transfer coefficient β are positive constants independent of the temperature. W.l.o.g. we set $\varrho c_p = 1$ in the analysis.
6. The viscosity parameters ϵ and γ are positive.

Remark 2. If the thermal strain \mathbf{t} fulfills [Assumption 1 item 4a](#) and satisfies

$$\mathbf{t}(\theta) = -M \quad \text{if } \theta \leq -M \quad \text{and} \quad \mathbf{t}(\theta) = M \quad \text{if } \theta \geq M$$

for some $M > 0$, then the product $\theta \mathbf{t}'(\theta)$ is Lipschitz continuous.

Next we introduce suitable function spaces for the weak formulation of [\(3\)–\(5\)](#).

Definition 3.

1. We define for $p \geq 2$ the (vector-valued) Sobolev space

$$\mathbf{W}_D^{1,p}(\Omega) := \{ \mathbf{u} \in \mathbf{W}^{1,p}(\Omega; \mathbb{R}^3) : \mathbf{u} = \mathbf{0} \text{ on } \Gamma_D \}.$$

2. We denote the dual space of $\mathbf{W}_D^{1,p}(\Omega)$ by $\mathbf{W}_D^{-1,p'}(\Omega)$, where $1/p + 1/p' = 1$.
3. We define for $p \geq 2$ the (matrix-valued) Lebesgue space

$$\mathbf{Q}^p(\Omega) := \{ \mathbf{q} \in \mathbf{L}^p(\Omega; \mathbb{R}_{trace}^{3 \times 3}) \}.$$

The following regularities for the initial conditions and the mechanical and thermal loads are assumed.

Assumption 4. Let $p, q \geq 2$ be fixed and define

$$v(p) \begin{cases} = 3p/(6-p) & \text{if } p < 6 \\ \in (\frac{3p}{3+p}, \infty) \text{ arbitrary} & \text{if } p \geq 6. \end{cases} \quad (6)$$

1. The initial conditions \mathbf{u}_0 , \mathbf{p}_0 and θ_0 have regularity

$$\mathbf{u}_0 \in \mathbf{W}_D^{1,p}(\Omega), \quad \mathbf{p}_0 \in \mathbf{Q}^p(\Omega) \quad \text{and} \quad \theta_0 \in W^{1,v(p)}(\Omega).$$

2. The volume and boundary loads ℓ and r belongs to the spaces

$$\ell \in L^q(0, T; \mathbf{W}_D^{-1,p}(\Omega)) \quad \text{and} \quad r \in L^{\frac{q}{2}}(0, T; W_{\diamond}^{-1,v(p)}(\Omega)).$$

Remark 5. The distinction of cases in the definition of $v(p)$ is due to the Sobolev embedding $L^{\frac{p}{2}}(\Omega) \hookrightarrow W_{\diamond}^{-1,v(p)}(\Omega)$ which becomes saturated for $p \geq 6$.

While conditions (1)–(2) are considered in the pointwise sense, the weak formulation of (3)–(5) is given by the following conditions, holding for almost all $t \in (0, T)$:

viscoplastic flow rule:

$$\begin{aligned} & \epsilon \int_{\Omega} \dot{\mathbf{p}} : (\mathbf{q} - \dot{\mathbf{p}}) \, d\mathbf{x} - \int_{\Omega} (\boldsymbol{\sigma} + \boldsymbol{\chi}) : (\mathbf{q} - \dot{\mathbf{p}}) \, d\mathbf{x} \\ & + \int_{\Omega} D(\mathbf{q}, \theta) \, d\mathbf{x} - \int_{\Omega} D(\dot{\mathbf{p}}, \theta) \, d\mathbf{x} \geq 0 \quad \text{for all } \mathbf{q} \in \mathbf{Q}^p(\Omega), \end{aligned} \quad (3')$$

balance of momentum:

$$\int_{\Omega} (\boldsymbol{\sigma} + \gamma \boldsymbol{\varepsilon}(\dot{\mathbf{u}})) : \boldsymbol{\varepsilon}(\mathbf{v}) \, d\mathbf{x} = \langle \boldsymbol{\ell}, \mathbf{v} \rangle \quad \text{for all } \mathbf{v} \in \mathbf{W}_D^{1,p'}(\Omega), \quad (4')$$

heat equation:

$$\begin{aligned} & \int_{\Omega} \dot{\theta} z \, d\mathbf{x} + \int_{\Omega} \kappa \nabla \theta \cdot \nabla z \, d\mathbf{x} + \int_{\Gamma} \beta \theta z \, ds \\ & = \langle r, z \rangle + \int_{\Omega} (\boldsymbol{\sigma} + \boldsymbol{\chi}) : \dot{\mathbf{p}} z \, d\mathbf{x} - \int_{\Omega} \theta \mathbf{t}'(\theta) : \mathbb{C}(\boldsymbol{\varepsilon}(\dot{\mathbf{u}}) - \dot{\mathbf{p}}) z \, d\mathbf{x} \\ & + \gamma \int_{\Omega} \boldsymbol{\varepsilon}(\dot{\mathbf{u}}) : \boldsymbol{\varepsilon}(\dot{\mathbf{u}}) z \, d\mathbf{x} \quad \text{for all } z \in L^{\frac{q}{q-2}}(0, T; W^{1,v(p)'}(\Omega)). \end{aligned} \quad (5')$$

We remark that $L^{\frac{q}{q-2}}(0, T; W^{1,v(p)'}(\Omega))$ is the dual space to $L^{\frac{q}{2}}(0, T; W_{\diamond}^{-1,v(p)}(\Omega))'$. Note that the balance of momentum (3) is equipped with mixed boundary conditions

$$\mathbf{u} = \mathbf{0} \quad \text{on } \Gamma_D \quad \text{and} \quad (\gamma \boldsymbol{\varepsilon}(\dot{\mathbf{u}}) + \boldsymbol{\sigma}) \mathbf{n} = \mathbf{s} \quad \text{on } \Gamma_N,$$

where \mathbf{n} is the outwards unit normal of Ω . The surface traction forces \mathbf{s} , together with volume loads, are summarized in $\boldsymbol{\ell}$. Moreover, the heat equation (5) is endowed with Robin boundary conditions whose right hand side enters r .

Finally, we present the assumptions on the domain.

Assumption 6.

1. $\Omega \subset \mathbb{R}^3$ is a bounded domain with Lipschitz boundary Γ , see, e.g., (Grisvard, 1985, Definition 1.2.1.1). The boundary Γ is divided into disjoint measurable parts Γ_N and Γ_D such that $\Gamma = \Gamma_N \dot{\cup} \Gamma_D$. Furthermore, Γ_N is an open and Γ_D is a closed subset of Γ with positive measure.
2. The set $\Omega \cup \Gamma_N$ is regular in the sense of Gröger (1989), which will be necessary to obtain $\mathbf{W}^{1,p}$ regularity (for some $p > 2$) of a solution of (4), as well as for the following assumption on maximal parabolic regularity.
3. In addition, the domain Ω is assumed to be smooth enough such that the operator related to (8) satisfies maximal parabolic regularity in $W_{\diamond}^{-1,v(p)}(\Omega)$; for a precise definition see Definition 33.

Remark 7.

1. In 3D, there is no simple characterization for [Assumption 6 item 2](#); cf. ([Haller-Dintelmann et al., 2009](#), Theorem 5.4). For example $\Omega \cup \Gamma_N$ is regular in the sense of Gröger if $\Omega \subset \mathbb{R}^3$ is a Lipschitzian polyhedron and $\bar{\Gamma}_N \cap \Gamma_D$ is a finite union of line segments; see ([Haller-Dintelmann et al., 2009](#), Corollary 5.5).
2. [Assumption 6 item 3](#) is not very restrictive because there exists $\hat{v} > 2$ such that the operator related to (8) satisfies maximal parabolic regularity in $W_\diamond^{-1,v(p)}(\Omega)$ for $\hat{v}' \leq v(p) \leq \hat{v}$ (where \hat{v}' is the conjugate exponent of \hat{v}); cf. [Lemma 35](#) and [Lemma 36](#).

3. Existence and Uniqueness of a Solution to the Model

In this section we prove the existence and uniqueness of the solution to the thermoviscoplastic system (1)–(5). We start with a precise definition of a solution and re-iterate our main theorem followed by a detailed roadmap of its proof. The major part of this section is a rigorous proof of the main theorem.

Theorem 8 (Main Theorem). *Suppose that [Assumption 1](#) and [Assumption 6](#) hold. There exists $\bar{p} > 2$ such that for all $2 < p \leq \bar{p}$, there exists $\bar{q} > 2$ (depending on p) such that for all $\bar{q} \leq q < \infty$ and right hand sides (ℓ, r) and initial conditions $(\mathbf{u}_0, \mathbf{p}_0, \theta_0)$ as in [Assumption 4](#), there exists a unique weak solution $(\mathbf{u}, \mathbf{p}, \theta, \boldsymbol{\sigma}, \boldsymbol{\chi})$ of (1)–(5) such that*

$$\begin{aligned} \mathbf{u} &\in W^{1,q}(0, T; \mathbf{W}_D^{1,p}(\Omega)), & \mathbf{p} &\in W^{1,q}(0, T; \mathbf{Q}^p(\Omega)), \\ \boldsymbol{\sigma} &\in W^{1,q}(0, T; \mathbf{L}^p(\Omega)), & \boldsymbol{\chi} &\in W^{1,q}(0, T; \mathbf{L}^p(\Omega)), \\ \theta &\in W^{1,\frac{q}{2}}(0, T; W_\diamond^{-1,v(p)}(\Omega)) \cap L^{\frac{q}{2}}(0, T; W^{1,v(p)}(\Omega)). \end{aligned}$$

That is, (1)–(2) hold almost everywhere in $\Omega \times (0, T)$ and (3')–(5') hold almost everywhere in $(0, T)$, and the initial conditions $\mathbf{u}(0) = \mathbf{u}_0$, $\mathbf{p}(0) = \mathbf{p}_0$ and $\theta(0) = \theta_0$ are satisfied.

For simplicity, we will refer to (3') in the sequel as (3) and similarly for (4) and (5) but always have in mind the weak form of the respective equation.

Next we present the roadmap of the proof. The conditions required for the indices p and q will be collected in the course of the proof. Throughout, capital Greek letters refer to solution operators of certain equations.

1. We first consider (1)–(4) for a fixed temperature field $\theta \in L^1(0, T; L^1(\Omega))$ and prove the existence of a unique solution (\mathbf{u}, \mathbf{p}) . This gives rise to the definition of the solution map

$$\begin{aligned} L^1(0, T; L^1(\Omega)) \ni \theta \mapsto \Lambda(\theta) &:= (\mathbf{u}, \mathbf{p}, \boldsymbol{\sigma}, \boldsymbol{\chi}) \in W^{1,q}(0, T; \mathbf{W}_D^{1,p}(\Omega)) \times \\ &\times W^{1,q}(0, T; \mathbf{Q}^p(\Omega)) \times [W^{1,q}(0, T; \mathbf{L}^p(\Omega))]^2. \end{aligned} \quad (7)$$

Individual components of this map will be referred to as $\Lambda^u(\theta)$ etc. or simply $\mathbf{u}(\theta)$. The final result is given in [Proposition 13](#).

2. The results of [item 1](#) naturally lead to the definition of a reduced problem for the temperature alone. To show the existence of a unique solution, we apply Banach's fixed point theorem, which requires a number of preparatory steps.

- (a) In order to apply maximal parabolic regularity results, we split the temperature field $\theta = \vartheta + \vartheta_{\text{init}}$ into its homogeneous and inhomogeneous parts w.r.t. the initial conditions. They are defined by

$$\int_{\Omega} \dot{\vartheta}_{\text{init}} z \, d\mathbf{x} + \int_{\Omega} \kappa \nabla \vartheta_{\text{init}} \cdot \nabla z \, d\mathbf{x} + \int_{\Gamma} \beta \vartheta_{\text{init}} z \, ds = 0, \quad (8a)$$

$$\vartheta_{\text{init}}(0) = \theta_0 \quad (8b)$$

and

$$\begin{aligned} & \int_{\Omega} \dot{\vartheta} z \, d\mathbf{x} + \int_{\Omega} \kappa \nabla \vartheta \cdot \nabla z \, d\mathbf{x} + \int_{\Gamma} \beta \vartheta z \, ds \\ &= \langle r, z \rangle + \int_{\Omega} (\boldsymbol{\sigma}(\vartheta + \vartheta_{\text{init}}) + \boldsymbol{\chi}(\vartheta + \vartheta_{\text{init}})) : \dot{\mathbf{p}}(\vartheta + \vartheta_{\text{init}}) z \, d\mathbf{x} \\ &- \int_{\Omega} (\vartheta + \vartheta_{\text{init}}) \mathbf{t}'(\vartheta + \vartheta_{\text{init}}) : \mathbb{C}(\boldsymbol{\varepsilon}(\dot{\mathbf{u}}(\vartheta + \vartheta_{\text{init}})) - \dot{\mathbf{p}}(\vartheta + \vartheta_{\text{init}})) z \, d\mathbf{x} \\ &+ \gamma \int_{\Omega} \boldsymbol{\varepsilon}(\dot{\mathbf{u}}(\vartheta + \vartheta_{\text{init}})) : \boldsymbol{\varepsilon}(\dot{\mathbf{u}}(\vartheta + \vartheta_{\text{init}})) z \, d\mathbf{x}, \end{aligned} \quad (9a)$$

$$\vartheta(0) = 0 \quad (9b)$$

for all $z \in L^{\frac{q}{q-2}}(0, T; W^{1, v(p)'}(\Omega))$, respectively. By standard results (see [Lemma 32](#)), ϑ_{init} satisfies

$$\vartheta_{\text{init}} \in W^{1, \infty}(0, T; W_{\diamond}^{-1, v(p)}(\Omega)) \cap L^{\infty}(0, T; W^{1, v(p)}(\Omega)). \quad (10)$$

- (b) The right hand side of [\(9a\)](#), without the term involving r , defines a map

$$\begin{aligned} \mathcal{R} : L^q(0, T; L^p(\Omega)) &\rightarrow L^{\frac{q}{2}}(0, T; L^{\frac{p}{2}}(\Omega)), \\ \mathcal{R}(\vartheta) &:= (\boldsymbol{\sigma}(\vartheta + \vartheta_{\text{init}}) + \boldsymbol{\chi}(\vartheta + \vartheta_{\text{init}})) : \dot{\mathbf{p}}(\vartheta + \vartheta_{\text{init}}) \\ &- (\vartheta + \vartheta_{\text{init}}) \mathbf{t}'(\vartheta + \vartheta_{\text{init}}) : \mathbb{C}(\boldsymbol{\varepsilon}(\dot{\mathbf{u}}(\vartheta + \vartheta_{\text{init}})) - \dot{\mathbf{p}}(\vartheta + \vartheta_{\text{init}})) \\ &+ \gamma \boldsymbol{\varepsilon}(\dot{\mathbf{u}}(\vartheta + \vartheta_{\text{init}})) : \boldsymbol{\varepsilon}(\dot{\mathbf{u}}(\vartheta + \vartheta_{\text{init}})). \end{aligned}$$

In [Lemma 14](#) we prove the Lipschitz property of \mathcal{R} .

- (c) We next define the following three maps in order to construct the solution operator of the heat equation with right hand side $\mathcal{R}(\vartheta)$. To complete the right hand side of [\(9a\)](#), we define the affine map

$$\begin{aligned} \mathcal{F} : L^{\frac{q}{2}}(0, T; L^{\frac{p}{2}}(\Omega)) &\rightarrow L^{\frac{q}{2}}(0, T; W_{\diamond}^{-1, v(p)}(\Omega)), \\ \mathcal{F}(f) &:= f + r, \end{aligned}$$

composed of an embedding plus an addition of the thermal loads.

The second map

$$\begin{aligned}\Pi &: L^{\frac{q}{2}}(0, T; W_{\diamond}^{-1, v(p)}(\Omega)) \\ &\rightarrow W_0^{1, \frac{q}{2}}(0, T; W_{\diamond}^{-1, v(p)}(\Omega)) \cap L^{\frac{q}{2}}(0, T; W^{1, v(p)}(\Omega)), \\ \Pi(f) &:= \vartheta\end{aligned}$$

is linear and it is given by the unique solution of

$$\int_{\Omega} \dot{\vartheta} z \, d\mathbf{x} + \int_{\Omega} \kappa \nabla \vartheta \cdot \nabla z \, d\mathbf{x} + \int_{\Gamma} \beta \vartheta z \, ds = \langle f, z \rangle \quad (11)$$

for all $z \in L^{\frac{q}{q-2}}(0, T; W^{1, v(p)}(\Omega))$. Here we benefit from maximal parabolic regularity results; see [Lemma 16](#).

Finally, we denote by \mathcal{E} the compact embedding

$$\begin{aligned}\mathcal{E} &: W_0^{1, \frac{q}{2}}(0, T; W_{\diamond}^{-1, v(p)}(\Omega)) \cap L^{\frac{q}{2}}(0, T; W^{1, v(p)}(\Omega)) \\ &\hookrightarrow C([0, T]; L^p(\Omega)),\end{aligned}$$

see [Lemma 17](#), which imposes a lower bound on q .

- (d) In virtue of the above, we can define the reduced formulation of (1)–(5) in terms of the temperature alone as a fixed-point problem, $\vartheta = \Theta(\vartheta)$, where

$$\begin{aligned}\Theta &: L^{\infty}(0, T; L^p(\Omega)) \rightarrow L^{\infty}(0, T; L^p(\Omega)), \\ \Theta(\vartheta) &:= \mathcal{E} \Pi \mathcal{F}(\mathcal{R}(\vartheta)).\end{aligned}$$

We show in [Lemma 18](#) the Lipschitz continuity of Θ .

- (e) Unfortunately, Θ is not necessarily contractive when defined on the entire time interval $(0, T)$. We therefore split the time interval into smaller parts. The application of the concatenation technique is aggravated by the fact the Lipschitz constant of Θ depends on the initial condition and thus the lengths of the subintervals might degrade. We overcome this problem by considering (1)–(5) iteratively on a sequence of intervals $[T_{n-1}, T_n]$ of equal lengths and prepend the unique solution already established on $[0, T_{n-1}]$.
3. The fixed-point problem provides a unique solution $\vartheta \in L^{\infty}(0, T; L^p(\Omega))$. From there, a unique solution $(\mathbf{u}, \mathbf{p}, \theta)$ as in [Theorem 8](#) can be deduced.

The following three subsections are arranged according to the structure above.

3.1. Existence and Uniqueness of a Solution for Given Temperature Field

In order to prove the existence and uniqueness of a solution of (1)–(4) for a given temperature field, we reformulate (1)–(4) as an ODE and use the Picard-Lindelöf theorem, following [Paoli & Petrov \(2012\)](#). We start with two lemmas which help to rephrase the balance of momentum (4) and the plastic flow rule (3).

Lemma 9. *There exists $\hat{p} > 2$ such that for all $2 \leq p \leq \hat{p}$ and $\mathbf{F} \in \mathbf{W}_D^{-1,p}(\Omega)$, there exists a unique solution $\mathbf{u} \in \mathbf{W}_D^{1,p}(\Omega)$ of*

$$\int_{\Omega} \gamma \varepsilon(\mathbf{u}) : \varepsilon(\mathbf{v}) \, d\mathbf{x} = \langle \mathbf{F}, \mathbf{v} \rangle \quad \text{for all } \mathbf{v} \in \mathbf{W}_D^{1,p'}(\Omega). \quad (12)$$

The corresponding solution operator $\Phi^{\mathbf{u}} : \mathbf{W}_D^{-1,p}(\Omega) \rightarrow \mathbf{W}_D^{1,p}(\Omega)$, $\mathbf{F} \mapsto \mathbf{u}$ is linear and bounded and satisfies the following estimate

$$\|\mathbf{u}\|_{\mathbf{W}_D^{1,p}(\Omega)} = \|\Phi^{\mathbf{u}}(\mathbf{F})\|_{\mathbf{W}_D^{1,p}(\Omega)} \leq C \gamma^{-1} \|\mathbf{F}\|_{\mathbf{W}_D^{-1,p}(\Omega)}. \quad (13)$$

The Lipschitz constant $C \gamma^{-1}$ is independent of $p \in [2, \hat{p}]$.

Proof. The result follows immediately from (Herzog et al., 2011, Theorem 1.1) with $\mathbf{b}(\cdot, \varepsilon(\mathbf{u})) := \gamma \varepsilon(\mathbf{u})$. Assumption 6 item 2 is used here. \square

Remark 10. *In the sequel, we use as \mathbf{F}*

$$\langle \mathbf{F}(\ell, \mathbf{u}, \mathbf{p}, \theta), \mathbf{v} \rangle := \int_{\Omega} \ell \cdot \mathbf{v} \, d\mathbf{x} - \int_{\Omega} \mathbb{C}(\varepsilon(\mathbf{u}) - \mathbf{p} - \mathbf{t}(\theta)) : \varepsilon(\mathbf{v}) \, d\mathbf{x} \quad (14)$$

with $\ell \in \mathbf{W}_D^{-1,p}(\Omega)$, $\mathbf{u} \in \mathbf{W}_D^{1,p}(\Omega)$, $\mathbf{p} \in \mathbf{Q}^p(\Omega)$ and $\theta \in L^1(\Omega)$. Then $\mathbf{F}(\ell, \mathbf{u}, \mathbf{p}, \theta) \in \mathbf{W}_D^{-1,p}(\Omega)$ holds and (12) (with \mathbf{u} replaced by $\dot{\mathbf{u}}$) equals the balance of momentum (4) at some fixed point in time.

Next, to handle the plastic flow rule (3), let us consider the following variational inequality

$$\varepsilon \mathbf{a} : (\mathbf{q} - \mathbf{a}) + D(\mathbf{q}, \theta) - D(\mathbf{a}, \theta) \geq \mathbf{f} : (\mathbf{q} - \mathbf{a}) \quad \text{for all } \mathbf{q} \in \mathbb{R}_{\text{trace}}^{3 \times 3}. \quad (15)$$

and prove its solvability for every fixed right hand side $\mathbf{f} \in \mathbb{R}_{\text{sym}}^{3 \times 3'} \simeq \mathbb{R}_{\text{sym}}^{3 \times 3}$ and temperature $\theta \in \mathbb{R}$. Note that since $\mathbf{f} : \mathbf{q} = [\mathbf{f}]^D : \mathbf{q}$ for all $\mathbf{q} \in \mathbb{R}_{\text{trace}}^{3 \times 3}$, the solution will depend only on the deviatoric part $[\mathbf{f}]^D$ of the right hand side.

Lemma 11. *For every fixed temperature $\theta \in \mathbb{R}$ and right hand side $\mathbf{f} \in \mathbb{R}_{\text{sym}}^{3 \times 3}$, there exists a unique solution $\mathbf{a} \in \mathbb{R}_{\text{trace}}^{3 \times 3}$ of (15) and it fulfills the inequality*

$$\varepsilon |\mathbf{a}| \leq |\mathbf{f}|. \quad (16)$$

Furthermore, the solution operator $\Phi^{\mathbf{p}} : \mathbb{R} \times \mathbb{R}_{\text{sym}}^{3 \times 3} \rightarrow \mathbb{R}_{\text{trace}}^{3 \times 3}$, $(\theta, \mathbf{f}) \mapsto \mathbf{a}$ is Lipschitz continuous. More precisely,

$$|\Phi^{\mathbf{p}}(\theta_1, \mathbf{f}_1) - \Phi^{\mathbf{p}}(\theta_2, \mathbf{f}_2)| \leq \varepsilon^{-1} |\mathbf{f}_1 - \mathbf{f}_2| + \varepsilon^{-1} L_{\Phi^{\mathbf{p}}} |\theta_1 - \theta_2|, \quad (17)$$

where $L_{\Phi^{\mathbf{p}}}$ just depends on the Lipschitz constant of the yield stress function σ_0 .

Proof. Existence and uniqueness: We can use (Han & Reddy, 1999, Theorem 6.6) to obtain a unique solution for every right hand side $\mathbf{f} \in \mathbb{R}_{\text{sym}}^{3 \times 3}$ and fixed temperature θ , because $\mathbf{q} \mapsto D(\mathbf{q}, \theta) := \sqrt{2/3} \sigma_0(\theta) |\mathbf{q}|$ is proper, convex and lower semicontinuous.

Estimate: We choose $\mathbf{q} = \mathbf{0}$ in (15) and get

$$-\epsilon \langle \mathbf{a}, \mathbf{a} \rangle - D(\mathbf{a}, \theta) \geq -\langle \mathbf{f}, \mathbf{a} \rangle \quad \text{or} \quad \epsilon |\mathbf{a}|^2 \leq \langle \mathbf{f}, \mathbf{a} \rangle - D(\mathbf{a}, \theta) \leq \|\mathbf{f}\| |\mathbf{a}|,$$

where we used that σ_0 is positive.

Lipschitz continuity: We consider $\mathbf{a}_1 = \Phi^{\mathbf{p}}(\theta_1, \mathbf{f}_1)$ and $\mathbf{a}_2 = \Phi^{\mathbf{p}}(\theta_2, \mathbf{f}_2)$ and get from (15)

$$\begin{aligned} \epsilon \langle \mathbf{a}_1, \mathbf{q} - \mathbf{a}_1 \rangle + D(\mathbf{q}, \theta_1) - D(\mathbf{a}_1, \theta_1) &\geq \langle \mathbf{f}_1, \mathbf{q} - \mathbf{a}_1 \rangle \quad \text{for all } \mathbf{q} \in \mathbb{R}_{\text{trace}}^{3 \times 3}, \\ \epsilon \langle \mathbf{a}_2, \mathbf{q} - \mathbf{a}_2 \rangle + D(\mathbf{q}, \theta_2) - D(\mathbf{a}_2, \theta_2) &\geq \langle \mathbf{f}_2, \mathbf{q} - \mathbf{a}_2 \rangle \quad \text{for all } \mathbf{q} \in \mathbb{R}_{\text{trace}}^{3 \times 3}. \end{aligned}$$

We choose $\mathbf{q} = \mathbf{a}_2$ in the first inequality and $\mathbf{q} = \mathbf{a}_1$ in the second and add both inequalities:

$$\epsilon |\mathbf{a}_1 - \mathbf{a}_2|^2 \leq \langle \mathbf{f}_1 - \mathbf{f}_2, \mathbf{a}_1 - \mathbf{a}_2 \rangle + \sqrt{\frac{2}{3}} [\sigma_0(\theta_1) - \sigma_0(\theta_2)] [|\mathbf{a}_1| - |\mathbf{a}_2|].$$

Using the Cauchy-Schwarz inequality and the Lipschitz continuity of σ_0 we get

$$|\Phi^{\mathbf{p}}(\theta_1, \mathbf{f}_1) - \Phi^{\mathbf{p}}(\theta_2, \mathbf{f}_2)| = |\mathbf{a}_1 - \mathbf{a}_2| \leq \epsilon^{-1} |\mathbf{f}_1 - \mathbf{f}_2| + \epsilon^{-1} L_{\Phi^{\mathbf{p}}} |\theta_1 - \theta_2|.$$

□

Remark 12. The inequality (15) (with $\mathbf{a} = \dot{\mathbf{p}}$ and $\mathbf{f} = \boldsymbol{\sigma} + \boldsymbol{\chi}$) corresponds to the formulation of the plastic flow rule (3) for a certain point in time and space (t, \mathbf{x}) . To see this, substitute \mathbf{q} by $(\mathbf{q} - \dot{\mathbf{p}})\varphi + \dot{\mathbf{p}}$ with $\varphi \in C_0^\infty(\Omega)$ arbitrary such that $0 \leq \varphi \leq 1$ in (3). The fundamental lemma of calculus of variations then yields, for almost all $\mathbf{x} \in \Omega$,

$$\epsilon \langle \dot{\mathbf{p}}, \mathbf{q} - \dot{\mathbf{p}} \rangle + D(\mathbf{q}, \theta) - D(\dot{\mathbf{p}}, \theta) \geq \langle \boldsymbol{\sigma} + \boldsymbol{\chi}, \mathbf{q} - \dot{\mathbf{p}} \rangle \quad \text{for all } \mathbf{q} \in \mathbb{R}_{\text{trace}}^{3 \times 3}.$$

With the solution operators $\Phi^{\mathbf{u}}$ and $\Phi^{\mathbf{p}}$ at hand, we can now prove the existence and uniqueness of a solution of (1)–(4) for a given temperature field.

Proposition 13 (Existence and uniqueness for given temperature field). *Let $\theta \in L^1(0, T; L^1(\Omega))$, $\mathbf{u}_0 \in \mathbf{W}_D^{1,p}(\Omega)$, $\mathbf{p}_0 \in \mathbf{Q}^p(\Omega)$ and $\boldsymbol{\ell} \in L^q(0, T; \mathbf{W}_D^{-1,p}(\Omega))$ be given; cf. Assumption 4. Then there exists a unique solution*

$$(\mathbf{u}, \mathbf{p}, \boldsymbol{\sigma}, \boldsymbol{\chi}) \in W^{1,q}(0, T; \mathbf{W}_D^{1,p}(\Omega)) \times W^{1,q}(0, T; \mathbf{Q}^p(\Omega)) \times [W^{1,q}(0, T; \mathbf{L}^p(\Omega))]^2$$

of (1)–(4) with $p \in [2, \hat{p}]$ and $1 < q < \infty$ where $\hat{p} > 2$ is determined by Lemma 9. Furthermore, the solution operator $\Lambda : \theta \mapsto (\mathbf{u}, \mathbf{p}, \boldsymbol{\sigma}, \boldsymbol{\chi})$ fulfills the following two properties.

1. The solution operator $\Lambda|_{L^q(0, T; L^p(\Omega))}$ is Lipschitz continuous with Lipschitz constant $L_\Lambda(T, q, \gamma^{-1}, \epsilon^{-1})$.
2. The image of Λ is bounded by $C(T, q, \gamma^{-1}, \epsilon^{-1}, \mathbf{u}_0, \mathbf{p}_0)$ independently of the temperature θ .

Proof. Existence: We can rewrite the balance of momentum (4) and the plastic flow rule (3) by means of the solution operators Φ^u and Φ^p defined in Lemma 9 and Lemma 11, respectively, cf. Remark 10 and Remark 12. Therefore we obtain the Banach space-valued ODE system

$$\begin{pmatrix} \dot{u} \\ \dot{p} \end{pmatrix} = \begin{pmatrix} \Phi^u(F(\ell, u, p, \theta)) \\ \Phi^p(\theta, \Phi^\sigma(u, p, \theta) + \Phi^x(u, p, \theta)) \end{pmatrix} =: \Phi^{up}(u, p). \quad (18)$$

The maps Φ^σ and Φ^x are defined by the algebraic relations (1) and (2), respectively. Note that the right hand side is non-autonomous since ℓ and θ depend on time. It follows from (13) and the pointwise estimate of (16) that Φ^{up} maps $L^q(0, T; \mathbf{W}_D^{1,p}(\Omega)) \times L^q(0, T; \mathbf{Q}^p(\Omega))$ into itself.

To apply the Picard-Lindelöf theorem we show that Φ^{up} is Lipschitz continuous uniformly in time. More precisely, we show the estimate

$$\begin{aligned} \|\Phi^{up}(u_1, p_1) - \Phi^{up}(u_2, p_2)\|_{\mathbf{W}_D^{1,p}(\Omega) \times \mathbf{Q}^p(\Omega)} \\ \leq C \| (u_1 - u_2, p_1 - p_2) \|_{\mathbf{W}_D^{1,p}(\Omega) \times \mathbf{Q}^p(\Omega)} \end{aligned}$$

for all $u_1, u_2 \in \mathbf{W}_D^{1,p}(\Omega)$ and $p_1, p_2 \in \mathbf{Q}^p(\Omega)$, where C is independent of the time t . We calculate

$$\begin{aligned} \|\Phi^{up}(u_1, p_1) - \Phi^{up}(u_2, p_2)\|_{\mathbf{W}_D^{1,p}(\Omega) \times \mathbf{Q}^p(\Omega)} \\ = \|\Phi^u(F(\ell, u_1, p_1, \theta)) - \Phi^u(F(\ell, u_2, p_2, \theta))\|_{\mathbf{W}_D^{1,p}(\Omega)} \\ + \|\Phi^p(\theta, \Phi^\sigma(u_1, p_1, \theta) + \Phi^x(u_1, p_1, \theta)) \\ - \Phi^p(\theta, \Phi^\sigma(u_2, p_2, \theta) + \Phi^x(u_2, p_2, \theta))\|_{\mathbf{Q}^p(\Omega)} \\ \stackrel{(13),(17)}{\leq} C \gamma^{-1} \|\mathbb{C}(\varepsilon(u_1) - p_1) - \mathbb{C}(\varepsilon(u_2) - p_2)\|_{L^p(\Omega)} \\ + \epsilon^{-1} \|\Phi^\sigma(u_1, p_1, \theta) + \Phi^x(u_1, p_1, \theta) - \Phi^\sigma(u_2, p_2, \theta) - \Phi^x(u_2, p_2, \theta)\|_{L^p(\Omega)}. \end{aligned}$$

Now we use the properties of \mathbb{C}, \mathbb{H} (Assumption 1) and the definitions of Φ^σ and Φ^x to get

$$\begin{aligned} \|\Phi^{up}(u_1, p_1) - \Phi^{up}(u_2, p_2)\|_{\mathbf{W}_D^{1,p}(\Omega) \times \mathbf{Q}^p(\Omega)} \\ \leq C(\gamma^{-1} + \epsilon^{-1}) \| (u_1 - u_2, p_1 - p_2) \|_{\mathbf{W}_D^{1,p}(\Omega) \times \mathbf{Q}^p(\Omega)}. \end{aligned}$$

Therefore, we obtain from the Picard-Lindelöf theorem (see (Gajewski et al., 1974, Chapter V, Lemma 1.5) for the case $p = 2$) a unique solution $(\Lambda^u(\theta), \Lambda^p(\theta)) = (u(\theta), p(\theta)) \in W^{1,q}(0, T; \mathbf{W}_D^{1,p}(\Omega)) \times W^{1,q}(0, T; \mathbf{Q}^p(\Omega))$. For the remaining two components we set

$$\begin{aligned} \Lambda^\sigma(\theta) &:= \Phi^\sigma(u(\theta), p(\theta), \theta) = \mathbb{C}(\varepsilon(u(\theta)) - p(\theta) - t(\theta)), \\ \Lambda^x(\theta) &:= \Phi^x(u(\theta), p(\theta), \theta) = -\mathbb{H} p(\theta) \end{aligned}$$

and define the solution operator Λ of (1)–(4) as $\Lambda = (\Lambda^u, \Lambda^p, \Lambda^\sigma, \Lambda^x)$.

Lipschitz continuity: Let $\theta_i \in L^q(0, T; L^p(\Omega))$ and $(\mathbf{u}_i, \mathbf{p}_i, \boldsymbol{\sigma}_i, \boldsymbol{\chi}_i) := \Lambda(\theta_i)$ for $i = 1, 2$. First we integrate (18) and calculate with the same argument as above

$$\begin{aligned}
& \|(\mathbf{u}_1(t), \mathbf{p}_1(t)) - (\mathbf{u}_2(t), \mathbf{p}_2(t))\|_{\mathbf{W}_D^{1,p}(\Omega) \times \mathbf{Q}^p(\Omega)} \\
& \leq \int_0^t \|\Phi^{\mathbf{u}}(\mathbf{F}(\boldsymbol{\ell}, \mathbf{u}_1, \mathbf{p}_1, \theta_1)) - \Phi^{\mathbf{u}}(\mathbf{F}(\boldsymbol{\ell}, \mathbf{u}_2, \mathbf{p}_2, \theta_2))\|_{\mathbf{W}_D^{1,p}(\Omega)} \, ds \\
& \quad + \int_0^t \|\Phi^{\mathbf{p}}(\theta_1, \boldsymbol{\sigma}_1 + \boldsymbol{\chi}_1) - \Phi^{\mathbf{p}}(\theta_2, \boldsymbol{\sigma}_2 + \boldsymbol{\chi}_2)\|_{\mathbf{Q}^p(\Omega)} \, ds \\
& \leq C(\gamma^{-1} + \epsilon^{-1}) \int_0^t \|(\mathbf{u}_1, \mathbf{p}_1) - (\mathbf{u}_2, \mathbf{p}_2)\|_{\mathbf{W}_D^{1,p}(\Omega) \times \mathbf{Q}^p(\Omega)} \, ds \\
& \quad + C(\gamma^{-1} + \epsilon^{-1}) \int_0^t \|\theta_1 - \theta_2\|_{L^p(\Omega)} \, ds.
\end{aligned}$$

Next we obtain from Gronwall's lemma and Hölder's inequality

$$\begin{aligned}
& \|(\mathbf{u}_1(t), \mathbf{p}_1(t)) - (\mathbf{u}_2(t), \mathbf{p}_2(t))\|_{\mathbf{W}_D^{1,p}(\Omega) \times \mathbf{Q}^p(\Omega)} \\
& \leq C(\gamma^{-1} + \epsilon^{-1}) \int_0^t \|\theta_1 - \theta_2\|_{L^p(\Omega)} \, ds \, e^{C(\gamma^{-1} + \epsilon^{-1})t} \\
& \leq C(\gamma^{-1} + \epsilon^{-1}) \|\theta_1 - \theta_2\|_{L^q(0,t;L^p(\Omega))} t^{\frac{q-1}{q}} e^{C(\gamma^{-1} + \epsilon^{-1})t}.
\end{aligned}$$

Further, we infer again as above

$$\begin{aligned}
& \|(\dot{\mathbf{u}}_1(t), \dot{\mathbf{p}}_1(t)) - (\dot{\mathbf{u}}_2(t), \dot{\mathbf{p}}_2(t))\|_{\mathbf{W}_D^{1,p}(\Omega) \times \mathbf{Q}^p(\Omega)} \\
& \leq C(\gamma^{-1} + \epsilon^{-1}) \|(\mathbf{u}_1(t), \mathbf{p}_1(t)) - (\mathbf{u}_2(t), \mathbf{p}_2(t))\|_{\mathbf{W}_D^{1,p}(\Omega) \times \mathbf{Q}^p(\Omega)} \\
& \quad + C(\gamma^{-1} + \epsilon^{-1}) \|\theta_1(t) - \theta_2(t)\|_{L^p(\Omega)}
\end{aligned}$$

and together we conclude

$$\begin{aligned}
& \|(\Lambda^{\mathbf{u}}(\theta_1), \Lambda^{\mathbf{p}}(\theta_1)) - (\Lambda^{\mathbf{u}}(\theta_2), \Lambda^{\mathbf{p}}(\theta_2))\|_{W^{1,q}(0,T;\mathbf{W}_D^{1,p}(\Omega)) \times W^{1,q}(0,T;\mathbf{Q}^p(\Omega))} \\
& = \|(\mathbf{u}_1, \mathbf{p}_1) - (\mathbf{u}_2, \mathbf{p}_2)\|_{W^{1,q}(0,T;\mathbf{W}_D^{1,p}(\Omega)) \times W^{1,q}(0,T;\mathbf{Q}^p(\Omega))} \\
& \leq C(T, q, \gamma^{-1}, \epsilon^{-1}) \|\theta_1 - \theta_2\|_{L^q(0,T;L^p(\Omega))}.
\end{aligned}$$

The Lipschitz continuity of $\Lambda^{\boldsymbol{\sigma}}$ and $\Lambda^{\boldsymbol{\chi}}$ is clear.

Boundedness: Finally, we have to show that the image of Λ is a bounded. We prove this with the same techniques as above. Let $\theta \in L^1(0, T; L^1(\Omega))$ and $(\mathbf{u}, \mathbf{p}, \boldsymbol{\sigma}, \boldsymbol{\chi}) := \Lambda(\theta)$. First we integrate (18) and calculate

$$\begin{aligned}
& \|(\mathbf{u}(t), \mathbf{p}(t)) - (\mathbf{u}_0, \mathbf{p}_0)\|_{\mathbf{W}_D^{1,p}(\Omega) \times \mathbf{Q}^p(\Omega)} \\
& \leq \int_0^t \|\Phi^{\mathbf{u}}(\mathbf{F}(\boldsymbol{\ell}, \mathbf{u}, \mathbf{p}, \theta))\|_{\mathbf{W}_D^{1,p}(\Omega)} \, ds + \int_0^t \|\Phi^{\mathbf{p}}(\theta, \boldsymbol{\sigma} + \boldsymbol{\chi})\|_{\mathbf{Q}^p(\Omega)} \, ds \\
& \leq C(\gamma^{-1} + \epsilon^{-1}) \int_0^t \|(\mathbf{u}, \mathbf{p})\|_{\mathbf{W}_D^{1,p}(\Omega)} \, ds + C\gamma^{-1} \int_0^t \|\boldsymbol{\ell}\|_{\mathbf{W}_D^{-1,p}(\Omega)} \, ds
\end{aligned}$$

$$+ C(\gamma^{-1} + \epsilon^{-1})t,$$

where we used the estimates (13), (16) and the boundedness of the thermal strain \mathbf{t} . It follows from Gronwall's lemma that

$$\begin{aligned} & \|(\mathbf{u}(t), \mathbf{p}(t))\|_{\mathbf{W}_D^{1,p}(\Omega) \times \mathbf{Q}^p(\Omega)} \\ & \leq \left[C\gamma^{-1} \int_0^t \|\boldsymbol{\ell}\|_{\mathbf{W}_D^{-1,p}(\Omega)} ds + C(\gamma^{-1} + \epsilon^{-1})t \right] e^{C(\gamma^{-1} + \epsilon^{-1})t} \\ & \quad + \|(\mathbf{u}_0, \mathbf{p}_0)\|_{\mathbf{W}_D^{1,p}(\Omega) \times \mathbf{Q}^p(\Omega)} e^{C(\gamma^{-1} + \epsilon^{-1})t} \end{aligned}$$

holds. Therefore, we conclude that (\mathbf{u}, \mathbf{p}) is bounded in $L^\infty(0, T; \mathbf{W}_D^{1,p}(\Omega)) \times L^\infty(0, T; \mathbf{Q}^p(\Omega))$ independently of $\theta \in L^1(0, T; L^1(\Omega))$. In the second step we use again (18) and calculate for almost all $t \in (0, T)$ with the same techniques as above

$$\begin{aligned} \|(\dot{\mathbf{u}}(t), \dot{\mathbf{p}}(t))\|_{\mathbf{W}_D^{1,p}(\Omega) \times \mathbf{Q}^p(\Omega)} & \leq C(\gamma^{-1} + \epsilon^{-1}) \|(\mathbf{u}(t), \mathbf{p}(t))\|_{\mathbf{W}_D^{1,p}(\Omega) \times \mathbf{Q}^p(\Omega)} \\ & \quad + C\gamma^{-1} \|\boldsymbol{\ell}(t)\|_{\mathbf{W}_D^{-1,p}(\Omega)} + C(\gamma^{-1} + \epsilon^{-1}). \end{aligned}$$

Now we conclude that $(\dot{\mathbf{u}}, \dot{\mathbf{p}})$ is bounded in $L^q(0, T; \mathbf{W}_D^{1,p}(\Omega)) \times L^q(0, T; \mathbf{Q}^p(\Omega))$ independently of $\theta \in L^1(0, T; L^1(\Omega))$ and together with the first step we obtain the boundedness of $(\Lambda^{\mathbf{u}}(\theta), \Lambda^{\mathbf{p}}(\theta)) = (\mathbf{u}(\theta), \mathbf{p}(\theta))$ in $W^{1,q}(0, T; \mathbf{W}_D^{1,p}(\Omega)) \times W^{1,q}(0, T; \mathbf{Q}^p(\Omega))$. The boundedness of Λ^σ and Λ^χ is clear. \square

3.2. Results for the Reduced Model

In this subsection we prove the Lipschitz continuity and the contractivity (on small time intervals) of the fixed point mapping Θ in order to apply the Banach fixed point theorem. A subsequent concatenation argument then yields a unique solution (1)–(5) on the entire time interval. As was already mentioned in the roadmap, the fixed point mapping $\Theta = \mathcal{E} \Pi \mathcal{F} \mathcal{R}$ is a combination of four individual mappings and therefore we start with proving some properties for these.

We begin with the right hand side \mathcal{R} of the homogeneous part of the temperature equation (5), see (9).

Lemma 14. *Suppose $2 < p \leq \hat{p}$ (determined by Lemma 9) and $2 < q < \infty$. Then the mapping $\mathcal{R} : L^q(0, T; L^p(\Omega)) \rightarrow L^{\frac{q}{2}}(0, T; L^{\frac{p}{2}}(\Omega))$, defined by*

$$\begin{aligned} \mathcal{R}(\vartheta) & := (\boldsymbol{\sigma}(\vartheta + \vartheta_{init}) + \boldsymbol{\chi}(\vartheta + \vartheta_{init})) : \dot{\mathbf{p}}(\vartheta + \vartheta_{init}) \\ & \quad - (\vartheta + \vartheta_{init}) \mathbf{t}'(\vartheta + \vartheta_{init}) : \mathbb{C}(\boldsymbol{\varepsilon}(\dot{\mathbf{u}}(\vartheta + \vartheta_{init})) - \dot{\mathbf{p}}(\vartheta + \vartheta_{init})) \\ & \quad + \gamma \boldsymbol{\varepsilon}(\dot{\mathbf{u}}(\vartheta + \vartheta_{init})) : \boldsymbol{\varepsilon}(\dot{\mathbf{u}}(\vartheta + \vartheta_{init})) \end{aligned}$$

is Lipschitz continuous with Lipschitz constant $L_{\mathcal{R}}(T, q, \gamma^{-1}, \epsilon^{-1}, \mathbf{u}_0, \mathbf{p}_0)$ and this Lipschitz constant does not increase when the interval length T shrinks.

Proof. Take $\vartheta_1, \vartheta_2 \in L^q(0, T; L^p(\Omega))$ and set $\theta_1 = \vartheta_1 + \vartheta_{\text{init}}$ and $\theta_2 = \vartheta_2 + \vartheta_{\text{init}}$ respectively. Notice that, according to (10) and Corollary 38, ϑ_{init} satisfies

$$\vartheta_{\text{init}} \in W^{1,\infty}(0, T; W_{\diamond}^{-1,v(p)}(\Omega)) \cap L^{\infty}(0, T; W^{1,v(p)}(\Omega)) \hookrightarrow C([0, T]; L^p(\Omega)).$$

We calculate

$$\begin{aligned} & \|\mathcal{R}(\vartheta_1) - \mathcal{R}(\vartheta_2)\|_{L^{\frac{q}{2}}(0, T; L^{\frac{p}{2}}(\Omega))} \\ & \leq \|(\sigma(\theta_1) - \sigma(\theta_2) + \chi(\theta_1) - \chi(\theta_2))\|_{L^q(0, T; L^p(\Omega))} \|\dot{\mathbf{p}}(\theta_1)\|_{L^q(0, T; L^p(\Omega))} \\ & \quad + \|\sigma(\theta_2) + \chi(\theta_2)\|_{L^q(0, T; L^p(\Omega))} \|\dot{\mathbf{p}}(\theta_1) - \dot{\mathbf{p}}(\theta_2)\|_{L^q(0, T; L^p(\Omega))} \\ & \quad + \|\theta_1 \mathbf{t}'(\theta_1) - \theta_2 \mathbf{t}'(\theta_2)\|_{L^q(0, T; L^p(\Omega))} \|\mathbb{C}(\varepsilon(\dot{\mathbf{u}}(\theta_1)) - \dot{\mathbf{p}}(\theta_2))\|_{L^q(0, T; L^p(\Omega))} \\ & \quad + \|\theta_2 \mathbf{t}'(\theta_2)\|_{L^q(0, T; L^p(\Omega))} \|\mathbb{C}(\varepsilon(\dot{\mathbf{u}}(\theta_1)) - \varepsilon(\dot{\mathbf{u}}(\theta_2)))\|_{L^q(0, T; L^p(\Omega))} \\ & \quad + \|\theta_2 \mathbf{t}'(\theta_2)\|_{L^q(0, T; L^p(\Omega))} \|\mathbb{C}(\dot{\mathbf{p}}(\theta_1) - \dot{\mathbf{p}}(\theta_2))\|_{L^q(0, T; L^p(\Omega))} \\ & \quad + \gamma \|\varepsilon(\dot{\mathbf{u}}(\theta_1)) - \varepsilon(\dot{\mathbf{u}}(\theta_2))\|_{L^q(0, T; L^p(\Omega))} \|\varepsilon(\dot{\mathbf{u}}(\theta_1))\|_{L^q(0, T; L^p(\Omega))} \\ & \quad + \gamma \|\varepsilon(\dot{\mathbf{u}}(\theta_2))\|_{L^q(0, T; L^p(\Omega))} \|\varepsilon(\dot{\mathbf{u}}(\theta_1)) - \varepsilon(\dot{\mathbf{u}}(\theta_2))\|_{L^q(0, T; L^p(\Omega))} \\ & \leq C(T, q, \gamma^{-1}, \epsilon^{-1}, \mathbf{u}_0, \mathbf{p}_0) \|\theta_1 - \theta_2\|_{L^q(0, T; L^p(\Omega))} \\ & =: L_{\mathcal{R}} \|\vartheta_1 - \vartheta_2\|_{L^q(0, T; L^p(\Omega))}. \end{aligned}$$

Here we have used the Lipschitz continuity and boundedness of the solution operator Λ (see Proposition 13) and of the mapping $\theta \mathbf{t}'(\theta)$ (see Assumption 1). Moreover, the proof of Proposition 13 shows that the Lipschitz constant of Λ increases monotonically with T . \square

Furthermore, we have for the other required mappings the following properties, which are easy to verify.

Lemma 15. *Suppose $2 < p, q < \infty$. Then the affine mapping*

$$\mathcal{F} : L^{\frac{q}{2}}(0, T; L^{\frac{p}{2}}(\Omega)) \rightarrow L^{\frac{q}{2}}(0, T; W_{\diamond}^{-1,v(p)}(\Omega)), \quad f \mapsto f + r$$

(defined via the embedding) is Lipschitz continuous with some Lipschitz constant $L_{\mathcal{F}}$ independent of T .

Proof. see Remark 5 \square

Next we consider the Lipschitz continuity of the solution operator Π of the heat equation (11) with general right hand side. Notice that we benefit from maximal parabolic regularity results at this point.

Lemma 16. *Suppose $2 < p < \infty$ such that $v(p) \leq \hat{v}$ (determined by Lemma 35) and $2 < q < \infty$ hold. Then the solution operator*

$$\begin{aligned} \Pi : L^{\frac{q}{2}}(0, T; W_{\diamond}^{-1,v(p)}(\Omega)) & \rightarrow W_0^{1,\frac{q}{2}}(0, T; W_{\diamond}^{-1,v(p)}(\Omega)) \cap L^{\frac{q}{2}}(0, T; W^{1,v(p)}(\Omega)) \\ & f \mapsto \vartheta \end{aligned}$$

related to (11) is linear and bounded, i.e., it satisfies the following estimate,

$$\|\vartheta\|_{W_0^{1,\frac{q}{2}}(0,T;W_\diamond^{-1,v(p)}(\Omega)) \cap L^{\frac{q}{2}}(0,T;W^{1,v(p)}(\Omega))} \leq L_\Pi \|f\|_{L^{\frac{q}{2}}(0,T;W_\diamond^{-1,v(p)}(\Omega))}. \quad (19)$$

The Lipschitz constant L_Π does not increase when the interval length T shrinks.

Proof. For the first statement we benefit from [Assumption 6 item 3](#); cf. [Remark 7](#). Notice that [Assumption 6 item 3](#) is satisfied in the case $\hat{v}' \leq v(p) \leq \hat{v}$ anyway. Only in the case $\frac{3}{2} < v(p) < \hat{v}' < 2$ does it constitute an additional assumption. The estimate follows easily from the closed graph theorem since the operator A related to (11) is closed. Consider now a function f in $L^{\frac{q}{2}}(0,T;W_\diamond^{-1,v(p)}(\Omega))$ and $T' < T$ and define the shifted (right-aligned) extension by zero,

$$(If)(t) := \begin{cases} 0 & \text{for } 0 \leq t \leq T - T', \\ f(t - (T - T')) & \text{for } T - T' < t \leq T. \end{cases}$$

The second result follows using the identity $\Pi If = I \Pi_{[0,T']} f$ where $\Pi_{[0,T']}$ is the restriction of Π to the time interval $[0, T']$; compare ([Hömborg et al., 2009/10](#), Lemma 3.16 (i)). \square

The next result concerns the embedding into spaces of continuous functions in time.

Lemma 17. *Suppose $2 < p < \infty$ and $2 < q$ (depending on p) sufficiently large. The embedding*

$$\mathcal{E} : W_0^{1,\frac{q}{2}}(0,T;W_\diamond^{-1,v(p)}(\Omega)) \cap L^{\frac{q}{2}}(0,T;W^{1,v(p)}(\Omega)) \rightarrow C([0,T];L^p(\Omega)), \quad f \mapsto f$$

is continuous. The Lipschitz constant $L_\mathcal{E}$ does not increase when the interval length T shrinks.

Proof. We refer to [Corollary 38](#) for the embedding and the precise link between p and q . Consider now a function f in $W_0^{1,\frac{q}{2}}(0,T';W_\diamond^{-1,v(p)}(\Omega)) \cap L^{\frac{q}{2}}(0,T';W^{1,v(p)}(\Omega))$ with $T' < T$ and define the shifted extension by zero I as in the proof of [Lemma 16](#). The second result follows as above, compare ([Hömborg et al., 2009/10](#), Lemma 3.16 (ii)). \square

With these lemmas we are now able to prove the Lipschitz continuity of the fixed point operator Θ .

Lemma 18. *Suppose $2 < p \leq \hat{p}$ (determined by [Lemma 9](#)) such that $v(p) \leq \hat{v}$ (determined by [Lemma 35](#)) and $2 < q < \infty$ (depending on p) sufficiently large. Then the mapping $\Theta : L^\infty(0,T;L^p(\Omega)) \rightarrow L^\infty(0,T;L^p(\Omega))$, $\Theta = \mathcal{E} \Pi \mathcal{F} \mathcal{R}$, is Lipschitz continuous, and it satisfies*

$$\|\Theta(\vartheta_1) - \Theta(\vartheta_2)\|_{L^\infty(0,T;L^p(\Omega))} \leq L_\Theta(q, \gamma^{-1}, \epsilon^{-1}, \mathbf{u}_0, \mathbf{p}_0) T^{\frac{1}{q}} \|\vartheta_1 - \vartheta_2\|_{L^\infty(0,T;L^p(\Omega))}$$

for all $\vartheta_1, \vartheta_2 \in L^\infty(0,T;L^p(\Omega))$. Hence the Lipschitz constant becomes arbitrarily small for sufficiently small $T > 0$.

Proof. Choose $\vartheta_1, \vartheta_2 \in L^\infty(0, T; L^p(\Omega))$. We use [Lemma 14](#) through [Lemma 17](#) and the Hölder inequality to obtain the following estimate.

$$\begin{aligned} \|\Theta(\vartheta_1) - \Theta(\vartheta_2)\|_{L^\infty(0, T; L^p(\Omega))} &= \|\mathcal{E} \Pi \mathcal{F}(\mathcal{R}(\vartheta_1)) - \mathcal{E} \Pi \mathcal{F}(\mathcal{R}(\vartheta_2))\|_{L^\infty(0, T; L^p(\Omega))} \\ &\leq L_{\mathcal{E}} L_{\Pi} L_{\mathcal{F}} L_{\mathcal{R}} \|\vartheta_1 - \vartheta_2\|_{L^q(0, T; L^p(\Omega))} \\ &\leq \underbrace{L_{\mathcal{E}} L_{\Pi} L_{\mathcal{F}} L_{\mathcal{R}}}_{=: L_{\Theta}} T^{\frac{1}{q}} \|\vartheta_1 - \vartheta_2\|_{L^\infty(0, T; L^p(\Omega))} \end{aligned}$$

with $2 < q < \infty$ sufficiently large such that the embedding \mathcal{E} is valid, cf. [Corollary 38](#). Notice that L_{Θ} can be chosen independently of T for all T bounded above by some constant, compare [Lemma 14](#), [Lemma 16](#) and [Lemma 17](#). \square

Finally, [Lemma 18](#), together with the observation that the Lipschitz constants for \mathcal{E} , Π , \mathcal{F} and \mathcal{R} do not increase when the interval length shrinks, results in the following corollary.

Corollary 19. *Suppose $2 < p \leq \hat{p}$ (determined by [Lemma 9](#)) such that $v(p) \leq \hat{v}$ (determined by [Lemma 35](#)). Then the mapping $\Theta : L^\infty(0, T_1; L^p(\Omega)) \rightarrow L^\infty(0, T_1; L^p(\Omega))$, $\Theta = \mathcal{E} \Pi \mathcal{F} \mathcal{R}$, is contractive for T_1 sufficiently small.*

The Banach fixed point theorem, together with a careful concatenation argument, shows the main result of [item 2](#) of the roadmap.

Proposition 20. *Suppose $2 < p \leq \hat{p}$ (determined by [Lemma 9](#)) such that $v(p) \leq \hat{v}$ (determined by [Lemma 35](#)) holds. Then the mapping*

$$\Theta : L^\infty(0, T; L^p(\Omega)) \rightarrow L^\infty(0, T; L^p(\Omega)), \quad \Theta = \mathcal{E} \Pi \mathcal{F} \mathcal{R}$$

has a unique fixed point.

Proof. We begin by fixing a value of q satisfying the conditions of [Lemma 18](#).

Existence of a fixed point on a small time interval $[0, T_1]$: We restrict the operator Θ to the time interval $[0, T_1]$ according to [Corollary 19](#) and denote it by $\Theta_{[0, T_1]}$. The analysis above is unaffected, and in particular we may use the same Lipschitz constants. Therefore we obtain, by means of the Banach fixed point theorem, a unique fixed point $\vartheta_{[0, T_1]} \in L^\infty(0, T_1; L^p(\Omega))$ of $\Theta_{[0, T_1]}$.

Concatenation argument: We split the time interval $[0, T]$ into N parts of equal length T_1 and define $T_n := nT_1$ for $n = 1, \dots, N$ where $T_N = T$. (It is clear that T/T_1 can be made integer by slightly reducing T_1 if necessary.) Analogously to the step above we denote by $\Theta_{[0, T_n]}$ the restriction of Θ to the time interval $[0, T_n]$. The same Lipschitz constants are still valid for $\Theta_{[0, T_n]}$.

We use an induction argument to conclude the existence of a unique fixed point $\vartheta_{[0, T_n]}$ for $\Theta_{[0, T_n]}$, provided that the existence of a unique fixed point $\vartheta_{[0, T_{n-1}]}$ for $\Theta_{[0, T_{n-1}]}$ has already been established. In the following we denote by $f * g$ the concatenation of the functions f and g defined on neighboring time intervals.

Let $\vartheta_{[0, T_{n-1}]}$ be the unique fixed point for $\Theta_{[0, T_{n-1}]}$. Consider the mapping

$$L^\infty(T_{n-1}, T_n; L^p(\Omega)) \rightarrow L^\infty(0, T_n; L^p(\Omega)), \quad f \mapsto \Theta_{[0, T_n]}(\vartheta_{[0, T_{n-1}]} * f).$$

This mapping is contractive because we obtain, for $f_1, f_2 \in L^\infty(T_{n-1}, T_n; L^p(\Omega))$, with calculations similar as in the proof of [Lemma 18](#), the estimate

$$\begin{aligned}
& \|\Theta_{[0, T_n]}(\vartheta_{[0, T_{n-1}]} * f_1) - \Theta_{[0, T_n]}(\vartheta_{[0, T_{n-1}]} * f_2)\|_{L^\infty(0, T_n; L^p(\Omega))} \\
& \leq L_{\mathcal{E}} L_{\Pi} L_{\mathcal{F}} L_{\mathcal{R}} \|\vartheta_{[0, T_{n-1}]} * f_1 - \vartheta_{[0, T_{n-1}]} * f_2\|_{L^q(0, T_n; L^p(\Omega))} \\
& = L_{\mathcal{E}} L_{\Pi} L_{\mathcal{F}} L_{\mathcal{R}} \|f_1 - f_2\|_{L^q(T_{n-1}, T_n; L^p(\Omega))} \\
& \leq \underbrace{L_{\mathcal{E}} L_{\Pi} L_{\mathcal{F}} L_{\mathcal{R}} T_1^q}_{< 1 \text{ by assumption}} \|f_1 - f_2\|_{L^\infty(T_{n-1}, T_n; L^p(\Omega))}.
\end{aligned}$$

Therefore this auxiliary mapping has a unique fixed point $f \in L^\infty(T_{n-1}, T_n; L^p(\Omega))$ and we define $\vartheta_{[0, T_n]} := \vartheta_{[0, T_{n-1}]} * f \in L^\infty(0, T_n; L^p(\Omega))$.

It remains to show that $\vartheta_{[0, T_n]}$, obtained by concatenation, is indeed the unique fixed point of $\Theta_{[0, T_n]}$. Using the induction hypothesis and the result above, we find

$$\begin{aligned}
\Theta_{[0, T_n]}(\vartheta_{[0, T_n]}) &= \begin{cases} \Theta_{[0, T_{n-1}]}(\vartheta_{[0, T_{n-1}]}) & \text{for } t \in [0, T_{n-1}], \\ \Theta_{[0, T_n]}(\vartheta_{[0, T_{n-1}]} * f)|_{[T_{n-1}, T_n]} & \text{for } t \in [T_{n-1}, T_n] \end{cases} \\
&= \begin{cases} \vartheta_{[0, T_{n-1}]} & \text{for } t \in [0, T_{n-1}], \\ f & \text{for } t \in [T_{n-1}, T_n]. \end{cases}
\end{aligned}$$

In the first equality we used that $\Theta_{[0, T_n]}$ is an extension of $\Theta_{[0, T_{n-1}]}$ in the sense that $\Theta_{[0, T_n]}(f) \equiv \Theta_{[0, T_{n-1}]}(g)$ on $[0, T_{n-1}]$ holds, provided that $f \equiv g$ on $[0, T_{n-1}]$. This shows that $\vartheta_{[0, T_n]}$ is a fixed point of $\Theta_{[0, T_n]}$. The uniqueness follows from the uniqueness on both subintervals, and the induction step is complete.

When $n = N$ is reached, the assertion is proved since $\Theta = \Theta_{[0, T_N]}$. \square

3.3. Proof of the Main Theorem

Now we are in the position to prove our main [Theorem 8](#).

of [Theorem 8](#). First we set $\bar{p} := \max\{p \leq \hat{p} : v(p) \leq \hat{v}\}$, where \hat{p} and \hat{v} are determined by [Lemma 9](#) and [Lemma 35](#), respectively. We collect the results proven in [subsection 3.1](#) and [subsection 3.2](#) in order to show [item 3](#) in the roadmap given at the beginning of [section 3](#).

So far (at the end of [item 2e](#)) in the roadmap, we have established the existence of a unique fixed point ϑ of $\Theta = \mathcal{E} \Pi \mathcal{F} \mathcal{R}$. Leaving out the embedding \mathcal{E} , we obtain $\vartheta = \Pi \mathcal{F}(\mathcal{R}(\vartheta))$, which ensures the desired regularity for the homogeneous part of the temperature. Finally, we use [Proposition 13](#) to define the unique solution in the following way,

$$\begin{aligned}
& (\mathbf{u}, \mathbf{p}, \theta, \sigma, \chi) \\
& := (\Lambda^{\mathbf{u}}(\vartheta + \vartheta_{\text{init}}), \Lambda^{\mathbf{p}}(\vartheta + \vartheta_{\text{init}}), \vartheta + \vartheta_{\text{init}}, \Lambda^{\sigma}(\vartheta + \vartheta_{\text{init}}), \Lambda^{\chi}(\vartheta + \vartheta_{\text{init}})) \\
& \in W^{1,q}(0, T; \mathbf{W}_D^{1,p}(\Omega)) \times W^{1,q}(0, T; \mathbf{Q}^p(\Omega)) \\
& \quad \times W^{1, \frac{q}{2}}(0, T; W_{\diamond}^{-1, v(p)}(\Omega)) \cap L^{\frac{q}{2}}(0, T; W^{1, v(p)}(\Omega))
\end{aligned}$$

$$\times W^{1,q}(0, T; \mathbf{L}^p(\Omega)) \times W^{1,q}(0, T; \mathbf{L}^p(\Omega)).$$

Note that by (8), ϑ_{init} has the regularity required. \square

Remark 21 (Bounds for \bar{p} and \bar{q}).

1. The spatial p -integrability of the displacement \mathbf{u} and plastic strain \mathbf{p} is limited by \bar{p} . This follows from (Herzog et al., 2011, Theorem 1.1), which was used to prove Proposition 13, together with Lemma 35, which was needed to ensure maximal parabolic regularity for the operator related to the heat equation (9).
2. The q -integrability in time of the displacement \mathbf{u} and the plastic strain \mathbf{p} has to be larger than \bar{q} (in dependence of p) to ensure that the embedding \mathcal{E} is valid. Corollary 38 gives the precise link between p and \bar{q} as follows. Fix $p > 2$ and $v(p)$ by (6), then choose \bar{q} as

$$\begin{aligned} \text{(a) for } p < 6: \quad \bar{q} &> \frac{2}{a} \quad \text{with} \quad 0 < a < \begin{cases} 1 - \frac{3}{2p} & \text{if } p < 3 \\ \frac{1}{2} & \text{otherwise,} \end{cases} \\ \text{(b) for } p \geq 6: \quad \bar{q} &> \frac{2}{a} \quad \text{with} \quad 0 < a < \begin{cases} 1 - \frac{3}{2v(p)} + \frac{3}{2p} & \text{if } v(p) < p \\ \frac{1}{2} & \text{otherwise.} \end{cases} \end{aligned}$$

Furthermore, we obtain the following property for the solution operator of (1)–(5) which will be essential in proving the existence of a global minimizer in section 4.

Lemma 22 (Boundedness of the solution operator). *Under the assumptions of Theorem 8, the mapping $(\ell, r) \mapsto (\mathbf{u}, \mathbf{p}, \theta, \boldsymbol{\sigma}, \boldsymbol{\chi})$ from $L^q(0, T; \mathbf{W}_D^{-1,p}(\Omega)) \times L^{\frac{q}{2}}(0, T; W_\diamond^{-1,v(p)}(\Omega))$ into the spaces for $(\mathbf{u}, \mathbf{p}, \theta, \boldsymbol{\sigma}, \boldsymbol{\chi})$ as in Theorem 8 is bounded, i.e., the images of bounded sets are bounded.*

Proof. Suppose $\mathbf{B} \subset L^q(0, T; \mathbf{W}_D^{-1,p}(\Omega)) \times L^{\frac{q}{2}}(0, T; W_\diamond^{-1,v(p)}(\Omega))$ is a bounded set. Consider the image $(\mathbf{u}(\mathbf{B}), \mathbf{p}(\mathbf{B}), \theta(\mathbf{B}), \boldsymbol{\sigma}(\mathbf{B}), \boldsymbol{\chi}(\mathbf{B}))$. The proof of Proposition 13 shows that $\mathbf{u}(\mathbf{B})$, $\mathbf{p}(\mathbf{B})$, $\boldsymbol{\sigma}(\mathbf{B})$ and $\boldsymbol{\chi}(\mathbf{B})$ are bounded.

The boundedness of the temperatures $\theta(\mathbf{B}) = \vartheta(\mathbf{B}) + \vartheta_{\text{init}}$ can be shown using first the embedding according to Lemma 37, then estimate (19) and finally Gronwall's lemma. We choose $(\ell, r) \in \mathbf{B}$ and calculate

$$\begin{aligned} \|\vartheta(t)\|_{L^p(\Omega)} &\leq \|\vartheta\|_{L^\infty(0,t;L^p(\Omega))} \leq C \|\vartheta\|_{W_0^{1,\frac{q}{2}}(0,t;W_\diamond^{-1,v(p)}(\Omega)) \cap L^{\frac{q}{2}}(0,t;W^{1,v(p)}(\Omega))} \\ &\leq C \|f\|_{L^{\frac{q}{2}}(0,t;W_\diamond^{-1,v(p)}(\Omega))} \end{aligned}$$

where $f \in L^{\frac{q}{2}}(0, t; W_\diamond^{-1,v(p)}(\Omega))$ is defined as

$$\begin{aligned} \langle f, z \rangle &:= \langle r, z \rangle + \int_{\Omega} (\boldsymbol{\sigma}(\vartheta + \vartheta_{\text{init}}) + \boldsymbol{\chi}(\vartheta + \vartheta_{\text{init}})) : \dot{\mathbf{p}}(\vartheta + \vartheta_{\text{init}}) z \, d\mathbf{x} \\ &\quad - \int_{\Omega} (\vartheta + \vartheta_{\text{init}}) \mathbf{t}'(\vartheta + \vartheta_{\text{init}}) : \mathbb{C}(\boldsymbol{\varepsilon}(\dot{\mathbf{u}}(\vartheta + \vartheta_{\text{init}})) - \dot{\mathbf{p}}(\vartheta + \vartheta_{\text{init}})) z \, d\mathbf{x} \end{aligned}$$

$$+ \gamma \int_{\Omega} \varepsilon(\dot{\mathbf{u}}(\vartheta + \vartheta_{\text{init}})) : \varepsilon(\dot{\mathbf{u}}(\vartheta + \vartheta_{\text{init}})) z \, d\mathbf{x}$$

for $z \in L^{\frac{q}{q-2}}(0, t; W^{1, v(p)'}(\Omega))$. Since $\mathbf{u}(\mathbf{B})$ and $\mathbf{p}(\mathbf{B})$ are bounded in the desired spaces and ϑ_{init} is fixed with regularity as in (10) we can estimate in the following way,

$$\|\vartheta(t)\|_{L^p(\Omega)} \leq C + C\|\vartheta\|_{L^q(0, t; L^p(\Omega))}.$$

By the convexity of $z \mapsto z^q$ for $z \geq 0$, we obtain the following estimate

$$\|\vartheta(t)\|_{L^p(\Omega)}^q \leq C + C \int_0^t \|\vartheta\|_{L^p(\Omega)}^q$$

and by Gronwall's lemma

$$\|\vartheta(t)\|_{L^p(\Omega)}^q \leq C \quad \text{for all } t \in [0, T].$$

This means that ϑ is bounded in $L^\infty(0, T; L^p(\Omega))$. Next we can again use the estimate (19) to show that ϑ is bounded in the desired space. Together with the regularity of ϑ_{init} (see (10)) we obtain the assertion. \square

4. Optimal Control Problem

In this section we present an optimal control problem governed by the thermoviscoplastic model (1)–(5), where the controls consist of boundary forces and surface tractions ℓ , and heat sources r . The aim is to prove the existence of a global minimizer by way of weak continuity of the control-to-state mapping; see Proposition 29. As before, p' and q' denote the conjugate indices of p and q , respectively.

Problem 23. *Find optimal controls*

$$\ell^* \in W^{1, q}(0, T; \mathbf{L}^p(\Omega)), \quad r^* \in L^{\frac{q}{2}}(0, T; L^{\frac{p}{2}}(\Omega))$$

and corresponding states

$$\begin{aligned} \mathbf{u}^* &\in W^{1, q}(0, T; \mathbf{W}_D^{1, p}(\Omega)), & \mathbf{p}^* &\in W^{1, q}(0, T; \mathbf{Q}^p(\Omega)), \\ \theta^* &\in W^{1, \frac{q}{2}}(0, T; W_{\diamond}^{-1, v(p)}(\Omega)) \cap L^{\frac{q}{2}}(0, T; W^{1, v(p)}(\Omega)) \end{aligned}$$

which minimize

$$F(\ell, r, \mathbf{u}, \mathbf{p}, \theta) := \psi(\mathbf{u}, \mathbf{p}, \theta) + \beta_1 \|\ell\|_{W^{1, q}(0, T; \mathbf{L}^p(\Omega))}^{b_1} + \beta_2 \|r\|_{L^{\frac{q}{2}}(0, T; L^{\frac{p}{2}}(\Omega))}^{b_2}$$

subject to (1)–(5).

The following assumptions are imposed.

Assumption 24.

1. The function $\psi : W^{1,2}(0, T; \mathbf{W}_D^{1,2}(\Omega)) \times L^2(0, T; \mathbf{Q}^2(\Omega)) \times L^2(0, T; L^2(\Omega)) \rightarrow \mathbb{R}$ is weakly sequentially lower semi-continuous and bounded from below.
2. Cost parameters β_1, β_2 are positive, i.e. $\beta_1, \beta_2 > 0$.
3. The exponents b_1 and b_2 satisfy $1 < b_1, b_2 < \infty$.

There are many possibilities to create a suitable objective. For instance it could be of interest to optimize the displacement, the residual stress or the plastic strain, see the following example.

Example 25 (Possible choices for ψ).

- Let $\tilde{\mathbf{u}}$ be a desired displacement. Then the term $\psi(\mathbf{u}) = \frac{1}{2} \int_{\Omega} |\mathbf{u}(T) - \tilde{\mathbf{u}}|^2 \, d\mathbf{x}$ is a classical tracking-type objective for the terminal displacement.
- The objective $\psi(\mathbf{u}, \mathbf{p}, \theta) = \frac{1}{2} \int_{\Omega} |\mathbb{C}(\boldsymbol{\varepsilon}(\mathbf{u}(T)) - \mathbf{p}(T) - \mathbf{t}(\theta(T)))|^2 \, d\mathbf{x}$ seeks to minimize the terminal residual stress.

Both examples are more meaningful when a cooling phase is appended to the end of the control horizon $[0, T]$, which is easily accounted for by bound constraints for the controls, cf. [Remark 28](#).

Theorem 26 (Existence of an optimal control). *Under the assumptions of [Theorem 8](#) and [Assumption 24](#), there exists at least one global minimizer $(\ell^*, r^*, \mathbf{u}^*, \mathbf{p}^*, \theta^*)$ of [Problem 23](#) such that*

$$\begin{aligned} \ell^* &\in W^{1,q}(0, T; \mathbf{L}^p(\Omega)), & r^* &\in L^{\frac{q}{2}}(0, T; L^{\frac{p}{2}}(\Omega)), \\ \mathbf{u}^* &\in W^{1,q}(0, T; \mathbf{W}_D^{1,p}(\Omega)), & \mathbf{p}^* &\in W^{1,q}(0, T; \mathbf{Q}^p(\Omega)), \\ \theta^* &\in W^{1,\frac{q}{2}}(0, T; W_{\diamond}^{-1,v(p)}(\Omega)) \cap L^{\frac{q}{2}}(0, T; W^{1,v(p)}(\Omega)). \end{aligned}$$

Remark 27. With [Theorem 8](#) at hand we can define the control-to-state mapping $\mathcal{G} : (\ell, r) \mapsto (\mathbf{u}, \mathbf{p}, \theta)$ for the thermoviscoplastic model [\(1\)–\(5\)](#).

of [Theorem 26](#). The proof follows standard arguments so we can be brief. First we use the control-to-state-map $\mathcal{G} : (\ell, r) \mapsto (\mathbf{u}, \mathbf{p}, \theta)$ to define the reduced functional $f(\ell, r) := F(\ell, r, \mathcal{G}(\ell, r))$. The reduced objective f is bounded from below by [Assumption 24](#), we get the existence of an infimum z ,

$$z := \inf f(\ell, r) \in \mathbb{R}.$$

Let $\{(\ell_n, r_n)\}_{n \in \mathbb{N}}$ be a minimizing sequence with $\lim_{n \rightarrow \infty} f(\ell_n, r_n) = z$. Because ψ is bounded from below and the cost parameters are positive, we get the following bound for the control

$$\|\ell_n\|_{W^{1,q}(0, T; \mathbf{L}^p(\Omega))} + \|r_n\|_{L^{\frac{q}{2}}(0, T; L^{\frac{p}{2}}(\Omega))} \leq C.$$

Therefore, there exists a control

$$(\ell^*, r^*) \in W^{1,q}(0, T; \mathbf{L}^p(\Omega)) \times L^{\frac{q}{2}}(0, T; L^{\frac{p}{2}}(\Omega))$$

and a subsequence (again denoted with n) such that for $n \rightarrow \infty$

$$\begin{aligned}\ell_n &\rightharpoonup \ell^* && \text{weakly in } W^{1,q}(0, T; \mathbf{L}^p(\Omega)), \\ r_n &\rightharpoonup r^* && \text{weakly in } L^{\frac{q}{2}}(0, T; L^{\frac{p}{2}}(\Omega)).\end{aligned}$$

The functional f is weakly sequentially lower semi-continuous, because \mathcal{G} is weakly continuous, see [Proposition 29](#), and ψ and the norms are weakly sequentially lower semi-continuous. Therefore we get

$$z = \lim_{n \rightarrow \infty} F(\ell_n, r_n, \mathcal{G}(\ell_n, r_n)) = \lim_{n \rightarrow \infty} f(\ell_n, r_n) \geq f(\ell^*, r^*) \geq z,$$

and (ℓ^*, r^*) is a global minimizer. \square

Remark 28 (Additional constraints and objectives). *Theorem 26 remains true when the controls (ℓ, r) are restricted to a convex closed subset of their respective spaces, as described for instance by pointwise bounds. Moreover, pointwise state constraints for the temperature can be imposed as well. These are not only important from an application point of view, but they also justify the cut-off property assumed for the temperature dependent thermal strain function \mathbf{t} , see [Assumption 1](#). Furthermore, the stresses $\boldsymbol{\sigma}$ and $\boldsymbol{\chi}$ can be also included in the objective.*

Weak continuity of the control-to-state mapping

In this section we provide the remaining proof of the control-to-state mapping's weak sequential continuity.

Proposition 29. *Under the assumptions of [Theorem 8](#), and provided that $\ell \in W^{1,q'}(0, T; \mathbf{L}^{p'}(\Omega))$ holds, the control-to-state mapping $(\ell, r) \mapsto (\mathbf{u}, \mathbf{p}, \theta)$ from $L^q(0, T; \mathbf{W}_D^{-1,p}(\Omega)) \cap W^{1,q'}(0, T; \mathbf{L}^{p'}(\Omega)) \times L^{\frac{q}{2}}(0, T; W_\diamond^{-1,v(p)}(\Omega))$ into the spaces for $(\mathbf{u}, \mathbf{p}, \theta)$ as in [Theorem 8](#), is weakly sequentially continuous.*

Remark 30 (Additional regularity for controls). *Notice that the controls in [Theorem 26](#) satisfy the additional regularity assumptions of [Proposition 29](#) according to the choice of the norms in the objective of our optimization [Proposition 23](#) and the embedding $W^{1,q}(0, T; \mathbf{L}^p(\Omega)) \hookrightarrow W^{1,q'}(0, T; \mathbf{L}^{p'}(\Omega))$ for any $p, q > 2$. Furthermore, note that for $p \leq 3$, the embedding $W^{1,q'}(0, T; \mathbf{L}^{p'}(\Omega)) \hookrightarrow L^q(0, T; \mathbf{W}_D^{-1,p}(\Omega))$ holds.*

At first glance it may seem surprising that the passage to the limit for weakly convergent sequences is possible for each of the equations (1)–(5). It turns out that the passage to the limit for the equations (4) and (3) can be easily done using a reformulation, cf. ([Han & Reddy, 1999](#), Section 7.2) or ([Bartels & Roubíček \(2008\)](#)), and that the second and third term of the right hand side of the heat equation,

$$\dot{\theta} - \operatorname{div}(\kappa \nabla \theta) = r + \gamma \boldsymbol{\varepsilon}(\dot{\mathbf{u}}) : \boldsymbol{\varepsilon}(\dot{\mathbf{u}}) + (\boldsymbol{\sigma} + \boldsymbol{\chi}) : \dot{\mathbf{p}} - \theta \mathbf{t}'(\theta) : \mathbb{C}(\boldsymbol{\varepsilon}(\dot{\mathbf{u}}) - \dot{\mathbf{p}}),$$

cause the most difficulties due to their nonlinearities. In order to handle them we would expect to need some terms strongly convergent in suitable spaces, such as for example $\varepsilon(\dot{\mathbf{u}})$. To overcome these difficulties, we adapt a technique from [Bartels & Roubíček \(2008\)](#), which encompasses a joint treatment of the two terms in question rather than considering the limits individually.

We begin with the reformulation of the balance of momentum (4) and plastic flow rule (3), see [Lemma 31](#), and subsequently give the proof of [Proposition 29](#). For that we define the following variational inequality for a given temperature $\theta \in L^1(0, T; L^1(\Omega))$,

$$\begin{aligned} J(\mathbf{u}, \mathbf{p}, \theta; \mathbf{v}, \mathbf{q}) := & \quad (20) \\ & \epsilon \int_0^T \int_{\Omega} \dot{\mathbf{p}} : (\mathbf{q} - \dot{\mathbf{p}}) \, d\mathbf{x} \, dt \\ & + \int_0^T \int_{\Omega} \mathbb{C} [\varepsilon(\mathbf{u}) - \mathbf{p} - \mathbf{t}(\theta)] : (\varepsilon(\mathbf{v}) - \mathbf{q} - (\varepsilon(\dot{\mathbf{u}}) - \dot{\mathbf{p}})) \, d\mathbf{x} \, dt \\ & + \int_0^T \int_{\Omega} \mathbb{H} \mathbf{p} : (\mathbf{q} - \dot{\mathbf{p}}) \, d\mathbf{x} \, dt + \gamma \int_0^T \int_{\Omega} \varepsilon(\dot{\mathbf{u}}) : (\varepsilon(\mathbf{v}) - \varepsilon(\dot{\mathbf{u}})) \, d\mathbf{x} \, dt \\ & + \int_0^T \int_{\Omega} D(\mathbf{q}, \theta) \, d\mathbf{x} \, dt - \int_0^T \int_{\Omega} D(\dot{\mathbf{p}}, \theta) \, d\mathbf{x} \, dt - \int_0^T \langle \ell, \mathbf{v} - \dot{\mathbf{u}} \rangle \, dt \geq 0 \end{aligned}$$

for $(\mathbf{v}, \mathbf{q}) \in L^{q'}(0, T; \mathbf{W}_D^{1,p'}(\Omega)) \times L^q(0, T; \mathbf{Q}^p(\Omega))$. This inequality is related to the thermoviscoplastic system in the following way.

Lemma 31. *Let $p, q \geq 2$, $\mathbf{u} \in W^{1,q}(0, T; \mathbf{W}_D^{1,p}(\Omega))$ and $\mathbf{p} \in W^{1,q}(0, T; \mathbf{Q}^p(\Omega))$. Then inequality (20) is equivalent to (1)–(4).*

Proof. “ \Leftarrow ” Insert (1) and (2) into (3) and (4), respectively. Now substitute \mathbf{v} by $\mathbf{v} - \dot{\mathbf{u}}$ in (4), add (3) and (4) and integrate over time to obtain (20).

“ \Rightarrow ” First choose $\mathbf{v} = \dot{\mathbf{u}}$ in (20) to get

$$\begin{aligned} & \epsilon \int_0^T \int_{\Omega} \dot{\mathbf{p}} : (\mathbf{q} - \dot{\mathbf{p}}) \, d\mathbf{x} \, dt - \int_0^T \int_{\Omega} \mathbb{C} [\varepsilon(\mathbf{u}) - \mathbf{p} - \mathbf{t}(\theta)] : (\mathbf{q} - \dot{\mathbf{p}}) \, d\mathbf{x} \, dt \quad (21) \\ & + \int_0^T \int_{\Omega} \mathbb{H} \mathbf{p} : (\mathbf{q} - \dot{\mathbf{p}}) \, d\mathbf{x} \, dt + \int_0^T \int_{\Omega} D(\mathbf{q}, \theta) \, d\mathbf{x} \, dt - \int_0^T \int_{\Omega} D(\dot{\mathbf{p}}, \theta) \, d\mathbf{x} \, dt \geq 0 \end{aligned}$$

for all $\mathbf{q} \in L^q(0, T; \mathbf{Q}^p(\Omega))$. Next choose $\mathbf{q} = \dot{\mathbf{p}}$ and substitute \mathbf{v} by $\pm \mathbf{v} + \dot{\mathbf{u}}$ in (20) to get

$$\begin{aligned} & \int_0^T \int_{\Omega} \mathbb{C} [\varepsilon(\mathbf{u}) - \mathbf{p} - \mathbf{t}(\theta)] : \varepsilon(\mathbf{v}) \, d\mathbf{x} \, dt + \gamma \int_0^T \int_{\Omega} \varepsilon(\dot{\mathbf{u}}) : \varepsilon(\mathbf{v}) \, d\mathbf{x} \, dt \\ & = \int_0^T \langle \ell, \mathbf{v} \rangle \, dt \quad \text{for all } \mathbf{v} \in L^{q'}(0, T; \mathbf{W}_D^{1,p'}(\Omega)). \quad (22) \end{aligned}$$

Finally, substitute \mathbf{q} by $(\mathbf{q} - \dot{\mathbf{p}})\varphi + \dot{\mathbf{p}}$ with $\varphi \in C_0^\infty(0, T)$ and $0 \leq \varphi \leq 1$ and \mathbf{v} by $\varphi \mathbf{v}$ with $\varphi \in C_0^\infty(0, T)$ in (21) and (22), respectively, and use the fundamental lemma of the calculus of variations to get (1)–(4). \square

Notice the structural advantages of this formulation. In the proof of [Proposition 29](#) we benefit from the quadratic structure of several terms since we can exploit the lower semicontinuity in $L^2(0, T; \mathbf{L}^2(\Omega))$ to handle them.

Proof. [of [Proposition 29](#)] Let us consider sequences $\{(\ell_n, r_n)\}$

$$\begin{aligned}\ell_n &\rightharpoonup \ell^* && \text{weakly in } W^{1,q'}(0, T; \mathbf{L}^{p'}(\Omega)), \\ \ell_n &\rightharpoonup \ell^* && \text{weakly in } L^q(0, T; \mathbf{W}_D^{-1,p}(\Omega)), \\ r_n &\rightharpoonup r^* && \text{weakly in } L^{\frac{q}{2}}(0, T; W_\diamond^{-1,v(p)}(\Omega))\end{aligned}$$

and define $(\mathbf{u}_n, \mathbf{p}_n, \theta_n) := \mathcal{G}(\ell_n, r_n)$. We have to show that

$$(\mathbf{u}_n, \mathbf{p}_n, \theta_n) = \mathcal{G}(\ell_n, r_n) \rightharpoonup \mathcal{G}(\ell^*, r^*) =: (\mathbf{u}^*, \mathbf{p}^*, \theta^*).$$

Definition of a candidate $(\mathbf{u}^*, \mathbf{p}^*, \theta^*)$: The displacements $\{\mathbf{u}_n\}$, the plastic strains $\{\mathbf{p}_n\}$ and the temperatures $\{\theta_n\}$ are, by [Lemma 22](#), bounded independently of n in the following sense:

$$\begin{aligned}\|\mathbf{u}_n\|_{W^{1,q}(0,T;\mathbf{W}_D^{1,p}(\Omega))} + \|\mathbf{p}_n\|_{W^{1,q}(0,T;\mathbf{Q}^p(\Omega))} &\leq C, \\ \|\theta_n\|_{W^{1,\frac{q}{2}}(0,T;W_\diamond^{-1,v(p)}(\Omega)) \cap L^{\frac{q}{2}}(0,T;W^{1,v(p)}(\Omega))} &\leq C.\end{aligned}$$

Therefore, there exist $\mathbf{u}^* \in W^{1,q}(0, T; \mathbf{W}_D^{1,p}(\Omega))$, $\mathbf{p}^* \in W^{1,q}(0, T; \mathbf{Q}^p(\Omega))$, and $\theta^* \in W^{1,\frac{q}{2}}(0, T; W_\diamond^{-1,v(p)}(\Omega)) \cap L^{\frac{q}{2}}(0, T; W^{1,v(p)}(\Omega))$ and a subsequence (denoted by n again) such that

$$\begin{aligned}\mathbf{u}_n &\rightharpoonup \mathbf{u}^* && \text{weakly in } W^{1,q}(0, T; \mathbf{W}_D^{1,p}(\Omega)), \\ \mathbf{p}_n &\rightharpoonup \mathbf{p}^* && \text{weakly in } W^{1,q}(0, T; \mathbf{Q}^p(\Omega)), \\ \theta_n &\rightharpoonup \theta^* && \text{weakly in } W^{1,\frac{q}{2}}(0, T; W_\diamond^{-1,v(p)}(\Omega)) \cap L^{\frac{q}{2}}(0, T; W^{1,v(p)}(\Omega)), \\ \theta_n &\rightarrow \theta^* && \text{strongly in } C([0, T]; L^p(\Omega)) \quad (\text{use } \text{Corollary 38}).\end{aligned}$$

Candidate is admissible, i.e., $\mathcal{G}(\ell^*, r^*) = (\mathbf{u}^*, \mathbf{p}^*, \theta^*)$: The idea is to show that $(\mathbf{u}^*, \mathbf{p}^*, \theta^*)$ fulfills the inequality [\(20\)](#) (which is equivalent to [\(1\)–\(4\)](#) by [Lemma 31](#)) and the heat equation [\(5\)](#). In order to do this we prove for $n \rightarrow \infty$ and for arbitrary $\mathbf{q} \in L^q(0, T; \mathbf{Q}^p(\Omega))$, $\mathbf{v} \in L^{q'}(0, T; \mathbf{W}^{1,p'}(\Omega))$ and $\varphi \in L^{\frac{q}{q-2}}(0, T; W^{1,v(p)}(\Omega))$ the following items. For brevity, we write $Q = \Omega \times (0, T)$.

1. $\lim_{n \rightarrow \infty} \int_Q \dot{\mathbf{p}}_n : \mathbf{q} \, d(\mathbf{x}, t) = \int_Q \dot{\mathbf{p}}^* : \mathbf{q} \, d(\mathbf{x}, t)$
2. $\liminf_{n \rightarrow \infty} \int_Q \dot{\mathbf{p}}_n : \dot{\mathbf{p}}_n \, d(\mathbf{x}, t) \geq \int_Q \dot{\mathbf{p}}^* : \dot{\mathbf{p}}^* \, d(\mathbf{x}, t)$
3. $\lim_{n \rightarrow \infty} \int_Q \mathbb{C}(\varepsilon(\mathbf{u}_n) - \mathbf{p}_n) : (\varepsilon(\mathbf{v}) - \mathbf{q}) \, d(\mathbf{x}, t) = \int_Q \mathbb{C}(\varepsilon(\mathbf{u}^*) - \mathbf{p}^*) : (\varepsilon(\mathbf{v}) - \mathbf{q}) \, d(\mathbf{x}, t)$
4. $\liminf_{n \rightarrow \infty} \int_Q \mathbb{C}(\varepsilon(\mathbf{u}_n) - \mathbf{p}_n) : (\varepsilon(\dot{\mathbf{u}}_n) - \dot{\mathbf{p}}_n) \, d(\mathbf{x}, t) \geq \int_Q \mathbb{C}(\varepsilon(\mathbf{u}^*) - \mathbf{p}^*) : (\varepsilon(\dot{\mathbf{u}}^*) - \dot{\mathbf{p}}^*) \, d(\mathbf{x}, t)$

5. $\lim_{n \rightarrow \infty} \int_Q \mathbb{C} \mathbf{t}(\theta_n) : (\varepsilon(\mathbf{v}) - \mathbf{q}) \, d(\mathbf{x}, t) = \int_Q \mathbb{C} \mathbf{t}(\theta^*) : (\varepsilon(\mathbf{v}) - \mathbf{q}) \, d(\mathbf{x}, t)$
6. $\lim_{n \rightarrow \infty} \int_Q \mathbb{C} \mathbf{t}(\theta_n) : (\varepsilon(\dot{\mathbf{u}}_n) - \dot{\mathbf{p}}_n) \, d(\mathbf{x}, t) = \int_Q \mathbb{C} \mathbf{t}(\theta^*) : (\varepsilon(\dot{\mathbf{u}}^*) - \dot{\mathbf{p}}^*) \, d(\mathbf{x}, t)$
7. $\lim_{n \rightarrow \infty} \int_Q \mathbb{H} \mathbf{p}_n : \mathbf{q} \, d(\mathbf{x}, t) = \int_Q \mathbb{H} \mathbf{p}^* : \mathbf{q} \, d(\mathbf{x}, t)$
8. $\liminf_{n \rightarrow \infty} \int_Q \mathbb{H} \mathbf{p}_n : \dot{\mathbf{p}} \, d(\mathbf{x}, t) \geq \int_Q \mathbb{H} \mathbf{p}^* : \dot{\mathbf{p}} \, d(\mathbf{x}, t)$
9. $\lim_{n \rightarrow \infty} \int_Q \varepsilon(\dot{\mathbf{u}}_n) : \varepsilon(\mathbf{v}) \, d(\mathbf{x}, t) = \int_Q \varepsilon(\dot{\mathbf{u}}^*) : \varepsilon(\mathbf{v}) \, d(\mathbf{x}, t)$
10. $\liminf_{n \rightarrow \infty} \int_Q \varepsilon(\dot{\mathbf{u}}) : \varepsilon(\dot{\mathbf{u}}) \, d(\mathbf{x}, t) \geq \int_Q \varepsilon(\dot{\mathbf{u}}^*) : \varepsilon(\dot{\mathbf{u}}^*) \, d(\mathbf{x}, t)$
11. $\lim_{n \rightarrow \infty} \int_Q D(\mathbf{q}, \theta_n) \, d(\mathbf{x}, t) = \int_Q D(\mathbf{q}, \theta) \, d(\mathbf{x}, t)$
12. $\liminf_{n \rightarrow \infty} \int_Q D(\dot{\mathbf{p}}_n, \theta_n) \, d(\mathbf{x}, t) \geq \int_Q D(\dot{\mathbf{p}}^*, \theta) \, d(\mathbf{x}, t)$
13. $\lim_{n \rightarrow \infty} \int_0^T \langle \ell_n, \mathbf{v} \rangle \, dt = \int_0^T \langle \ell^*, \mathbf{v} \rangle \, dt$
14. $\lim_{n \rightarrow \infty} \int_0^T \langle \ell_n, \dot{\mathbf{u}}_n \rangle \, dt = \int_0^T \langle \ell^*, \dot{\mathbf{u}}^* \rangle \, dt$
15. $\lim_{n \rightarrow \infty} \int_Q \dot{\theta}_n \varphi \, d(\mathbf{x}, t) = \int_Q \dot{\theta}^* \varphi \, d(\mathbf{x}, t)$
16. $\lim_{n \rightarrow \infty} \int_Q \operatorname{div}(\kappa \nabla \theta_n) \varphi \, d(\mathbf{x}, t) = \int_Q \operatorname{div}(\kappa \nabla \theta^*) \varphi \, d(\mathbf{x}, t)$
17. $\lim_{n \rightarrow \infty} \int_Q r_n \varphi \, d(\mathbf{x}, t) = \int_Q r^* \varphi \, d(\mathbf{x}, t)$
18. $\lim_{n \rightarrow \infty} \int_Q \theta_n \mathbf{t}'(\theta_n) : \mathbb{C}(\varepsilon(\dot{\mathbf{u}}_n) - \dot{\mathbf{p}}_n) \varphi \, d(\mathbf{x}, t) = \int_Q \theta \mathbf{t}'(\theta^*) : \mathbb{C}(\varepsilon(\dot{\mathbf{u}}^*) - \dot{\mathbf{p}}^*) \varphi \, d(\mathbf{x}, t)$
19. $\lim_{n \rightarrow \infty} \int_Q (\boldsymbol{\sigma}_n + \boldsymbol{\chi}_n) : \dot{\mathbf{p}}_n \varphi + \gamma \varepsilon(\dot{\mathbf{u}}_n) : \varepsilon(\dot{\mathbf{u}}_n) \varphi \, d(\mathbf{x}, t)$
 $= \int_Q (\boldsymbol{\sigma}^* + \boldsymbol{\chi}^*) : \dot{\mathbf{p}}^* \varphi + \gamma \varepsilon(\dot{\mathbf{u}}^*) : \varepsilon(\dot{\mathbf{u}}^*) \varphi \, d(\mathbf{x}, t),$
where $\boldsymbol{\sigma}_n := \mathbb{C}(\varepsilon(\mathbf{u}_n) - \mathbf{p}_n - \mathbf{t}(\theta_n))$, $\boldsymbol{\chi}_n := -\mathbb{H} \mathbf{p}_n$ and analogously $\boldsymbol{\sigma}^*$, $\boldsymbol{\chi}^*$ are defined.

From [item 1](#) to [item 14](#) we can conclude that $(\mathbf{u}^*, \mathbf{p}^*, \theta^*)$ fulfills [\(20\)](#). Indeed, since $(\mathbf{u}_n, \mathbf{p}_n, \theta_n)$ verifies [\(20\)](#) we can write this as $0 \leq J(\mathbf{u}_n, \mathbf{p}_n, \theta_n; \mathbf{v}, \mathbf{q})$ for all (\mathbf{v}, \mathbf{q}) . Now we use [item 1](#) to [item 14](#) to get

$$\begin{aligned} 0 &\leq \limsup_{n \rightarrow \infty} J(\mathbf{u}_n, \mathbf{p}_n, \theta_n; \mathbf{v}, \mathbf{q}) \\ &= -\liminf_{n \rightarrow \infty} -J(\mathbf{u}_n, \mathbf{p}_n, \theta_n; \mathbf{v}, \mathbf{q}) \leq J(\mathbf{u}^*, \mathbf{p}^*, \theta^*; \mathbf{v}, \mathbf{q}). \end{aligned}$$

Using [item 15](#) to [item 19](#) we conclude that $(\mathbf{u}^*, \mathbf{p}^*, \theta^*)$ fulfills the heat equation [\(5\)](#).

Let us prove the items above. First of all, [item 1](#), [item 3](#), [item 7](#), [item 9](#), [item 13](#), [item 15](#), [item 16](#) and [item 17](#) are clear. Concerning [item 2](#) (and similarly [item 10](#)) we use the weak sequentially lower semi-continuity of the norm in $L^2(0, T; \mathbf{L}^2(\Omega))$ to get

$$\liminf_{n \rightarrow \infty} \int_Q \dot{\mathbf{p}}_n : \dot{\mathbf{p}}_n \, d(\mathbf{x}, t) \geq \int_Q \dot{\mathbf{p}}^* : \dot{\mathbf{p}}^* \, d(\mathbf{x}, t).$$

Concerning [item 4](#) (and similarly [item 8](#)) we benefit from the property $\mathbf{u} : \dot{\mathbf{u}} = \frac{1}{2} \frac{d}{dt}(\mathbf{u} : \mathbf{u})$ and the weak sequential lower semi-continuity of the norm in $\mathbf{L}^2(\Omega)$ to calculate

$$\liminf_{n \rightarrow \infty} \int_Q \mathbb{C}(\varepsilon(\mathbf{u}_n) - \mathbf{p}_n) : (\varepsilon(\dot{\mathbf{u}}_n) - \dot{\mathbf{p}}_n) \, d(\mathbf{x}, t)$$

$$\begin{aligned}
&= \liminf_{n \rightarrow \infty} \frac{1}{2} \int_{\Omega} \mathbb{C}(\varepsilon(\mathbf{u}_n(T)) - \mathbf{p}_n(T)) : (\varepsilon(\mathbf{u}_n(T)) - \mathbf{p}_n(T)) \, d\mathbf{x} \\
&\quad - \frac{1}{2} \int_{\Omega} \mathbb{C}(\varepsilon(\mathbf{u}_n(0)) - \mathbf{p}_n(0)) : (\varepsilon(\mathbf{u}_n(0)) - \mathbf{p}_n(0)) \, d\mathbf{x} \\
&\geq \frac{1}{2} \int_{\Omega} \mathbb{C}(\varepsilon(\mathbf{u}^*(T)) - \mathbf{p}^*(T)) : (\varepsilon(\mathbf{u}^*(T)) - \mathbf{p}^*(T)) \, d\mathbf{x} \\
&\quad - \frac{1}{2} \int_{\Omega} \mathbb{C}(\varepsilon(\mathbf{u}_0^*) - \mathbf{p}_0^*) : (\varepsilon(\mathbf{u}_0^*) - \mathbf{p}_0^*) \, d\mathbf{x} \\
&= \int_Q \mathbb{C}(\varepsilon(\mathbf{u}^*) - \mathbf{p}^*) : (\varepsilon(\dot{\mathbf{u}}^*) - \dot{\mathbf{p}}^*) \, d(\mathbf{x}, t).
\end{aligned}$$

Notice that the weak convergence of the sequences $\{\varepsilon(\mathbf{u}_n(T))\}$ and $\{\mathbf{p}_n(T)\}$ in $\mathbf{L}^2(\Omega)$ follows from the continuity of the embedding $W^{1,q}(0, T; \mathbf{L}^p(\Omega))$ into $C([0, T]; \mathbf{L}^p(\Omega))$.

Concerning [item 6](#) (and similarly [item 5](#) and [item 18](#)) we calculate

$$\begin{aligned}
&\lim_{n \rightarrow \infty} \int_Q \mathbb{C} \mathbf{t}(\theta_n) : (\varepsilon(\dot{\mathbf{u}}_n) - \dot{\mathbf{p}}_n) \pm \mathbb{C} \mathbf{t}(\theta^*) : (\varepsilon(\dot{\mathbf{u}}_n) - \dot{\mathbf{p}}_n) \\
&\quad - \mathbb{C} \mathbf{t}(\theta^*) : (\varepsilon(\dot{\mathbf{u}}^*) - \dot{\mathbf{p}}^*) \, d(\mathbf{x}, t) \\
&\leq C \lim_{n \rightarrow \infty} \|\theta_n - \theta^*\|_{L^{q'}(0, T; L^{p'}(\Omega))} \|\varepsilon(\dot{\mathbf{u}}_n) - \dot{\mathbf{p}}_n\|_{L^q(0, T; \mathbf{L}^p(\Omega))} \\
&\quad + \int_Q \mathbb{C} \mathbf{t}(\theta^*) : (\varepsilon(\dot{\mathbf{u}}_n) - \varepsilon(\dot{\mathbf{u}}^*) - (\dot{\mathbf{p}}_n - \dot{\mathbf{p}}^*)) \, d(\mathbf{x}, t) = 0.
\end{aligned}$$

For the last term we use that $\mathbf{m} \mapsto \mathbb{C} \mathbf{t}(\theta^*) : \mathbf{m}$ with $L^p(0, T; \mathbf{L}^q(\Omega)) \rightarrow \mathbb{R}$ is a linear and continuous. Therefore, it is an element of $L^{p'}(0, T; \mathbf{L}^{q'}(\Omega))$ and we get with the weak convergence of $\varepsilon(\mathbf{u}_n)$ and \mathbf{p}_n that

$$\int_Q \mathbb{C} \mathbf{t}(\theta^*) : (\varepsilon(\dot{\mathbf{u}}_n) - \varepsilon(\dot{\mathbf{u}}^*) - (\dot{\mathbf{p}}_n - \dot{\mathbf{p}}^*)) \, d(\mathbf{x}, t) = 0.$$

Concerning [item 11](#) and [item 12](#) we can use the same arguments as above in combination with the Lipschitz continuity and positivity of σ_0 .

Concerning [item 14](#) we have the compact embedding $\mathbf{L}^{p'}(\Omega) \hookrightarrow \mathbf{W}_D^{-1, p'}(\Omega)$ and therefore the embedding $W^{1, q'}(0, T; \mathbf{L}^{p'}(\Omega)) \hookrightarrow L^{q'}(0, T; \mathbf{W}_D^{-1, p'}(\Omega))$ is also compact, see ([Simon, 1986](#), Theorem 3, (6.5)). That means that there exists a subsequence $\ell_n \rightarrow \ell^*$ in $L^{q'}(0, T; \mathbf{W}_D^{-1, p'}(\Omega))$ and

$$\begin{aligned}
&\lim_{n \rightarrow \infty} \int_0^T \langle \ell_n, \dot{\mathbf{u}}_n \rangle - \langle \ell^*, \dot{\mathbf{u}}^* \rangle \, dt \\
&\leq \lim_{n \rightarrow \infty} \|\ell_n - \ell^*\|_{L^{q'}(0, T; \mathbf{W}_D^{-1, p'}(\Omega))} \|\dot{\mathbf{u}}_n\|_{L^q(0, T; \mathbf{W}_D^{1, p}(\Omega))} \\
&\quad + \int_0^T \langle \ell^*, \dot{\mathbf{u}}_n - \dot{\mathbf{u}}^* \rangle \, dt = 0
\end{aligned}$$

follows.

Concerning [item 19](#) (following [\(Bartels & Roubíček, 2008, Proof of Proposition 4.6\)](#)) we test the plastic flow rule [\(3\)](#) by $\mathbf{q} = \mathbf{0}$ and $\mathbf{q} = 2\dot{\mathbf{p}}$ and get

$$D(\dot{\mathbf{p}}, \theta) = (\boldsymbol{\sigma} + \boldsymbol{\chi}) : \dot{\mathbf{p}} - \epsilon \dot{\mathbf{p}} : \dot{\mathbf{p}}.$$

Therefore, we can rephrase the term in [item 19](#) as

$$\begin{aligned} & \int_Q (\boldsymbol{\sigma}_n + \boldsymbol{\chi}_n) : \dot{\mathbf{p}}_n \varphi + \gamma \boldsymbol{\varepsilon}(\dot{\mathbf{u}}_n) : \boldsymbol{\varepsilon}(\dot{\mathbf{u}}_n) \varphi \, d(\mathbf{x}, t) \\ &= \int_Q D(\dot{\mathbf{p}}_n, \theta_n) \varphi + \epsilon \dot{\mathbf{p}}_n : \dot{\mathbf{p}}_n \varphi + \gamma \boldsymbol{\varepsilon}(\dot{\mathbf{u}}_n) : \boldsymbol{\varepsilon}(\dot{\mathbf{u}}_n) \varphi \, d(\mathbf{x}, t) \end{aligned}$$

Since $\{D(\dot{\mathbf{p}}_n, \theta_n)\}$ is bounded in $L^q(0, T; L^p(\Omega))$, there exists $\xi_1 \in L^q(0, T; L^p(\Omega))$ and a subsequence such that

$$D(\dot{\mathbf{p}}_n, \theta_n) \rightharpoonup \xi_1.$$

Similarly $\{\epsilon \dot{\mathbf{p}}_n : \dot{\mathbf{p}}_n\}$ and $\{\gamma \dot{\mathbf{u}}_n : \dot{\mathbf{u}}_n\}$ are bounded in $L^{\frac{p}{2}}(0, T; L^{\frac{q}{2}}(\Omega))$ and there exist $\xi_2, \xi_3 \in L^{\frac{p}{2}}(0, T; L^{\frac{q}{2}}(\Omega))$ and subsequences such that

$$\epsilon \dot{\mathbf{p}}_n : \dot{\mathbf{p}}_n \rightharpoonup \xi_2 \quad \text{and} \quad \gamma \dot{\mathbf{u}}_n : \dot{\mathbf{u}}_n \rightharpoonup \xi_3.$$

We use [item 12](#) and the weak sequential lower semicontinuity of the norm in $L^2(0, T; \mathbf{L}^2(\Omega))$ to calculate

$$\begin{aligned} & \int_0^T \int_{\Omega} \boldsymbol{\sigma}^* : \dot{\mathbf{p}}^* + \boldsymbol{\chi}^* : \dot{\mathbf{p}}^* + \gamma \boldsymbol{\varepsilon}(\dot{\mathbf{u}}^*) : \boldsymbol{\varepsilon}(\dot{\mathbf{u}}^*) \, d(\mathbf{x}, t) \\ &= \int_Q D(\theta^*, \dot{\mathbf{p}}^*) + \epsilon \dot{\mathbf{p}}^* : \dot{\mathbf{p}}^* + \gamma \boldsymbol{\varepsilon}(\dot{\mathbf{u}}^*) : \boldsymbol{\varepsilon}(\dot{\mathbf{u}}^*) \, d(\mathbf{x}, t) \\ &\leq \liminf_{n \rightarrow \infty} \int_Q D(\theta_n, \dot{\mathbf{p}}_n) + \epsilon \dot{\mathbf{p}}_n : \dot{\mathbf{p}}_n + \gamma \boldsymbol{\varepsilon}(\dot{\mathbf{u}}_n) : \boldsymbol{\varepsilon}(\dot{\mathbf{u}}_n) \, d(\mathbf{x}, t) \\ &\leq \limsup_{n \rightarrow \infty} \int_Q D(\theta_n, \dot{\mathbf{p}}_n) + \epsilon \dot{\mathbf{p}}_n : \dot{\mathbf{p}}_n + \gamma \boldsymbol{\varepsilon}(\dot{\mathbf{u}}_n) : \boldsymbol{\varepsilon}(\dot{\mathbf{u}}_n) \, d(\mathbf{x}, t) \\ &= \limsup_{n \rightarrow \infty} \int_Q \boldsymbol{\sigma}_n : \dot{\mathbf{p}}_n + \boldsymbol{\chi}_n : \dot{\mathbf{p}}_n + \gamma \boldsymbol{\varepsilon}(\dot{\mathbf{u}}_n) : \boldsymbol{\varepsilon}(\dot{\mathbf{u}}_n) \, d(\mathbf{x}, t) \\ &\quad - \int_Q (\boldsymbol{\sigma}_n + \gamma \boldsymbol{\varepsilon}(\dot{\mathbf{u}}_n)) : \boldsymbol{\varepsilon}(\dot{\mathbf{u}}_n) \, d(\mathbf{x}, t) + \int_0^T \langle \ell_n, \dot{\mathbf{u}}_n \rangle \, dt \\ &\quad \quad \quad [\text{by setting } \mathbf{v} = \dot{\mathbf{u}}_n \text{ in (4)}] \\ &= \limsup_{n \rightarrow \infty} \int_Q -\mathbb{C}(\boldsymbol{\varepsilon}(\mathbf{u}_n) - \mathbf{p}_n) : (\boldsymbol{\varepsilon}(\dot{\mathbf{u}}_n) - \dot{\mathbf{p}}_n) - \mathbb{H} \mathbf{p}_n : \dot{\mathbf{p}}_n \, d(\mathbf{x}, t) \\ &\quad + \int_Q \mathbb{C} \mathbf{t}(\theta_n) : (\boldsymbol{\varepsilon}(\dot{\mathbf{u}}_n) - \dot{\mathbf{p}}_n) \, d(\mathbf{x}, t) + \int_0^T \langle \ell_n, \dot{\mathbf{u}}_n \rangle \, dt \\ &\quad \quad \quad [\text{by using (1) and (2)}] \end{aligned}$$

$$\begin{aligned}
&= \int_Q -\mathbb{C}(\varepsilon(\mathbf{u}^*) - \mathbf{p}^*) : (\varepsilon(\dot{\mathbf{u}}^*) - \dot{\mathbf{p}}^*) - \mathbb{H}\mathbf{p}^* : \dot{\mathbf{p}}^* \, d(\mathbf{x}, t) \\
&\quad + \int_Q \mathbb{C}\mathbf{t}(\theta^*) : (\varepsilon(\dot{\mathbf{u}}^*) - \dot{\mathbf{p}}^*) \, d(\mathbf{x}, t) + \int_0^T \langle \ell^*, \dot{\mathbf{u}} \rangle \, dt \\
&\quad \quad \quad [\text{by using item 4, item 6, item 8 and item 14}] \\
&= \int_Q \boldsymbol{\sigma}^* : \dot{\mathbf{p}}^* + \boldsymbol{\chi}^* : \dot{\mathbf{p}}^* + \gamma \varepsilon(\dot{\mathbf{u}}^*) : \varepsilon(\dot{\mathbf{u}}^*) \, d(\mathbf{x}, t) \\
&\quad \quad \quad [\text{by setting } \mathbf{v} = \dot{\mathbf{u}}^* \text{ in (4)}].
\end{aligned}$$

Therefore, all inequalities are in fact equalities. Now we use that if

$$\lim_{n \rightarrow \infty} \int_Q a_n \, d\mathbf{x} = \int_Q a \, d\mathbf{x} \quad \text{with } a_n, a \geq 0,$$

then we get for arbitrary $\varphi \in L^\infty(0, T; L^\infty(\Omega))$

$$\lim_{n \rightarrow \infty} \int_Q (a_n - a) \varphi \, d\mathbf{x} \leq \text{ess sup } \varphi \cdot \lim_{n \rightarrow \infty} \int_Q (a_n - a) \, d\mathbf{x} = 0.$$

Hence we conclude that

$$\boldsymbol{\sigma}_n : \dot{\mathbf{p}}_n + \boldsymbol{\chi}_n : \dot{\mathbf{p}}_n + \gamma \dot{\mathbf{u}}_n : \dot{\mathbf{u}}_n \rightharpoonup \boldsymbol{\sigma}^* : \dot{\mathbf{p}}^* + \boldsymbol{\chi}^* : \dot{\mathbf{p}}^* + \gamma \dot{\mathbf{u}}^* : \dot{\mathbf{u}}^* \text{ in } L^1(0, T; L^1(\Omega)).$$

Since the weak limit is unique, we get

$$\boldsymbol{\sigma}_n : \dot{\mathbf{p}}_n + \boldsymbol{\chi}_n : \dot{\mathbf{p}}_n + \gamma \dot{\mathbf{u}}_n : \dot{\mathbf{u}}_n \rightharpoonup \boldsymbol{\sigma}^* : \dot{\mathbf{p}}^* + \boldsymbol{\chi}^* : \dot{\mathbf{p}}^* + \gamma \dot{\mathbf{u}}^* : \dot{\mathbf{u}}^* (= \xi_1 + \xi_2 + \xi_3)$$

in $L^{\frac{q}{2}}(0, T; W_\diamond^{-1, v(p)}(\Omega))$.

Convergence $(\mathbf{u}_n, \mathbf{p}_n, \theta_n) \rightharpoonup (\mathbf{u}^*, \mathbf{p}^*, \theta^*)$ for the entire sequence: In the step above we have shown that $(\mathbf{u}_n, \mathbf{p}_n, \theta_n) \rightharpoonup (\mathbf{u}^*, \mathbf{p}^*, \theta^*)$ for a subsequence. With the arguments above we can prove that every subsequence has a subsequence converging to $(\mathbf{u}^*, \mathbf{p}^*, \theta^*)$. Therefore, the entire sequence $(\mathbf{u}_n, \mathbf{p}_n, \theta_n)$ converges to $(\mathbf{u}^*, \mathbf{p}^*, \theta^*)$. □

Acknowledgment

The authors would like to thank Joachim Rehberg (WIAS Berlin) and Hannes Meinschmidt (TU Darmstadt) for fruitful discussions about maximal parabolic regularity theory.

Appendix A. Appendix

Appendix A.1. Semigroup theory

Lemma 32. *The solution ϑ_{init} of (8) satisfies*

$$\vartheta_{init} \in W^{1, \infty}(0, T; W_\diamond^{-1, v(p)}(\Omega)) \cap L^\infty(0, T; W^{1, v(p)}(\Omega)).$$

Proof. Notice, that the semigroup $(T(t))_t : W_\diamond^{-1,v(p)}(\Omega) \rightarrow W_\diamond^{-1,v(p)}(\Omega)$ with domain $W^{1,v(p)}$ of the operator $-A$ related to (8) is analytic and the solution is given by $\vartheta_{\text{init}}(t) = T(t)\theta_0$. We can estimate using the properties of an analytic semigroup in the following way,

$$\begin{aligned} \|-A\vartheta_{\text{init}}(t)\|_{W_\diamond^{-1,v(p)}(\Omega)} &= \|-AT(t)\theta_0\|_{W_\diamond^{-1,v(p)}(\Omega)} \\ &= \|T(t)(-A)\theta_0\|_{W_\diamond^{-1,v(p)}(\Omega)} \\ &\leq \|T(t)\|_{W_\diamond^{-1,v(p)}(\Omega) \rightarrow W_\diamond^{-1,v(p)}(\Omega)} \|-A\theta_0\|_{W_\diamond^{-1,v(p)}(\Omega)}. \end{aligned}$$

Using the equivalence of the graph norm $\|-A\cdot\|_{W_\diamond^{-1,v(p)}(\Omega)} + \|\cdot\|_{W_\diamond^{-1,v(p)}(\Omega)}$ and the norm of the space $W^{1,v(p)}(\Omega)$ we infer that

$$\begin{aligned} &\text{ess sup}_{t \in [0,T]} \|\vartheta_{\text{init}}(t)\|_{W^{1,v(p)}(\Omega)} \\ &\leq C \sup_{t \in [0,T]} \|T(t)\|_{W_\diamond^{-1,v(p)}(\Omega) \rightarrow W_\diamond^{-1,v(p)}(\Omega)} \|\theta_0\|_{W^{1,v(p)}(\Omega)} < C. \end{aligned}$$

In combination with $\dot{\vartheta}_{\text{init}} = -A\vartheta_{\text{init}} \in L^\infty(0,T; W_\diamond^{-1,v(p)}(\Omega))$ we obtain the regularity

$$\vartheta_{\text{init}} \in W^{1,\infty}(0,T; W_\diamond^{-1,v(p)}(\Omega)) \cap L^\infty(0,T; W^{1,v(p)}(\Omega)).$$

□

Appendix A.2. Maximal Parabolic Regularity

Definition 33 (Maximal parabolic regularity). *Let X be a Banach space and A a closed operator with dense domain $\mathcal{D} \subseteq X$. Suppose and $0 < T < \infty$. Then the operator A satisfies maximal parabolic regularity in X iff there exists $r \in (1, \infty)$ such that for any $f \in L^r(0,T; X)$ there is a unique function $w \in W_0^{1,r}(0,T; X) \cap L^r(0,T; \mathcal{D})$ which fulfills*

$$\dot{w} + Aw = f.$$

Remark 34. *It is well known that the property of maximal parabolic regularity of an operator A is independent of $r \in (1, \infty)$ and the choice of the time interval $[0, T]$; cf. (Dore, 1993, Theorem 4.2, Theorem 2.5).*

Lemma 35. *Under Assumption 6 item 2, there exists $\hat{v} > 2$ such that for every $v \in [2, \hat{v}]$ the operator related to (8) satisfies maximal parabolic regularity in $W_\diamond^{-1,v}(\Omega)$ independently of the time interval and the time integrability $r \in (1, \infty)$.*

Proof. See (Gröger, 1989, Theorem 1 and Remark 5) and Remark 34. □

Lemma 36. *Suppose A is a closed densely defined operator with domain \mathcal{D} satisfying maximal parabolic regularity in X . Then its adjoint operator A^* satisfies maximal parabolic regularity in \mathcal{D}' .*

Proof. By [Definition 33](#), an operator A satisfies maximal parabolic regularity in X iff there exists $r \in (1, \infty)$ such that the mapping

$$\partial_t + A : W_0^{1,r}(0, T; X) \cap L^r(0, T; \mathcal{D}) \rightarrow L^r(0, T; X)$$

is an isomorphism, where ∂_t denotes the weak time derivative. Then the adjoint operator is also an isomorphism

$$(\partial_t + A)^* : L^{r'}(0, T; X') \rightarrow (W_0^{1,r}(0, T; X) \cap L^r(0, T; \mathcal{D}))',$$

i.e. for all $g \in (W_0^{1,r}(0, T; X) \cap L^r(0, T; \mathcal{D}))'$, there exists a unique $\psi \in L^{r'}(0, T; X')$ such that

$$(\partial_t + A)^* \psi = \partial_t^* \psi + A^* \psi = g.$$

Furthermore, we obtain the following equation for given $g \in L^{r'}(0, T; \mathcal{D}') \subseteq (W_0^{1,r}(0, T; X) \cap L^r(0, T; \mathcal{D}))'$ and for all $\xi \in C_c^\infty(0, T; \mathcal{D})$ satisfying $\xi(t) = v(t)u$, where $v \in C_c^\infty(0, T)$ and $u \in \mathcal{D}$:

$$\int_0^T \langle (\partial_t + A)^* \psi, \xi \rangle_{\mathcal{D}} dt = \int_0^T \langle \partial_t^* \psi, \xi \rangle_{\mathcal{D}} dt + \int_0^T \langle A^* \psi, \xi \rangle_{\mathcal{D}} dt = \int_0^T \langle g, \xi \rangle_{\mathcal{D}} dt.$$

This means that

$$\begin{aligned} \langle u, \int_0^T v' \psi dt \rangle_{\mathcal{D}} &= \int_0^T \langle \psi, \partial_t \xi \rangle_{\mathcal{D}} dt = \int_0^T \langle \partial_t^* \psi, \xi \rangle_{\mathcal{D}} dt \\ &= \int_0^T \langle g - A^* \psi, \xi \rangle_{\mathcal{D}} dt \\ &= \langle u, \int_0^T (g - A^* \psi) v dt \rangle_{\mathcal{D}}. \end{aligned}$$

Since the equation above is satisfied for all $u \in \mathcal{D}$ we obtain

$$\int_0^T v' \psi dt = \int_0^T (g - A^* \psi) v dt$$

for all $v \in C_c^\infty(0, T)$, and the regularity of g implies that the distributional time derivative of ψ is regular and satisfies

$$-\partial_t \psi = g - A^* \psi \in L^{r'}(0, T; \mathcal{D}'). \quad (\text{A.1})$$

Therefore we have $\psi \in W^{1,r'}(0, T; \mathcal{D}') \cap L^{r'}(0, T; X')$. Now we use integration by parts ([Amann, 2005](#), Proposition 5.1) to get, for all $\varphi \in W_0^{1,r}(0, T; X) \cap L^r(0, T; \mathcal{D})$,

$$\int_0^T \langle g, \varphi \rangle_{\mathcal{D}} dt = \int_0^T \langle \partial_t^* \psi, \varphi \rangle_{\mathcal{D}} dt + \int_0^T \langle A^* \psi, \varphi \rangle_{\mathcal{D}} dt$$

$$\begin{aligned}
&= \int_0^T \langle \psi, \partial_t \varphi \rangle_X dt + \int_0^T \langle A^* \psi, \varphi \rangle_{\mathcal{D}} dt \\
&= \int_0^T \langle -\partial_t \psi, \varphi \rangle_{\mathcal{D}} dt + \int_0^T \langle \psi(T), \varphi(T) \rangle_{(X, \mathcal{D})_{1/r', r}} dt \\
&\quad + \int_0^T \langle A^* \psi, \varphi \rangle_{\mathcal{D}} dt,
\end{aligned}$$

where $(X, \mathcal{D})_{1/r', r}$ denotes the real interpolation space. Using (A.1) and the fact that φ was arbitrary, we obtain $\psi(T) = 0$ in $(X, \mathcal{D})'_{1/r', r} = (\mathcal{D}', X')_{1/r, r'} \hookrightarrow \mathcal{D}'$. Therefore for all $g \in L^{r'}(0, T; \mathcal{D}')$, there exists a unique $\psi \in W^{1, r'}(0, T; \mathcal{D}') \cap L^{r'}(0, T; X')$ such that

$$-\partial_t \psi + A^* \psi = g \quad \text{and} \quad \psi(T) = 0 \quad \text{in } \mathcal{D}'$$

hold. Finally we transform the time variable $s \rightarrow T - s$ and see that A^* satisfies maximal parabolic regularity in \mathcal{D}' . \square

Appendix A.3. Embeddings

We require the following results for embeddings in our analysis.

Lemma 37. Let $0 < a < \min \left\{ \frac{1}{2}, \frac{1}{2} - \frac{3}{2y} + \frac{3}{2z} \right\}$ and $\begin{cases} \frac{3z}{3+z} < y < \frac{3z}{3-z} & \text{if } p < 3 \\ \frac{3z}{3+z} < y < \infty, & \text{if } p = 3 \\ \frac{3z}{3+z} < y \leq \infty, & \text{otherwise.} \end{cases}$

1. For $aq < 2$ and $w < \frac{q}{2-aq}$ there is the compact embedding

$$W^{1, \frac{q}{2}}(0, T; W_{\diamond}^{-1, y}(\Omega)) \cap L^{\frac{q}{2}}(0, T; W^{1, y}(\Omega)) \hookrightarrow L^w(0, T; L^z(\Omega)).$$

2. For $aq > 2$ there is the compact embedding

$$W^{1, \frac{q}{2}}(0, T; W_{\diamond}^{-1, y}(\Omega)) \cap L^{\frac{q}{2}}(0, T; W^{1, y}(\Omega)) \hookrightarrow C([0, T]; L^z(\Omega)).$$

Proof. The embeddings follow with Corollary 8 of Simon (1986). Check all the assumptions therein by using Lemma 12 in Simon (1986). \square

Corollary 38. Fix $p > 2$.

1. Let $p < 6$ and thus $v(p) = 3p/(6-p)$; cf. (6).

Then for $q > \frac{2}{a}$ and $0 < a < \begin{cases} 1 - \frac{3}{2p} & \text{if } p < 3 \\ \frac{1}{2} & \text{otherwise} \end{cases}$ the following embeddings

are compact:

- (a) $W^{1, \frac{q}{2}}(0, T; W_{\diamond}^{-1, v(p)}(\Omega)) \cap L^{\frac{q}{2}}(0, T; W^{1, v(p)}(\Omega)) \hookrightarrow C([0, T]; L^p(\Omega)),$
- (b) $W_0^{1, \frac{q}{2}}(0, T; W_{\diamond}^{-1, v(p)}(\Omega)) \cap L^{\frac{q}{2}}(0, T; W^{1, v(p)}(\Omega)) \hookrightarrow C([0, T]; L^p(\Omega)).$

2. Let $p \geq 6$ and thus $v(p) \in (\frac{3p}{3+p}, \infty)$; cf. (6).

Then for $q > \frac{2}{a}$ and $0 < a < \begin{cases} 1 - \frac{3}{2v(p)} + \frac{3}{2p} & \text{if } v(p) < p \\ \frac{1}{2} & \text{otherwise} \end{cases}$ the following embeddings are compact:

- (a) $W^{1,\frac{q}{2}}(0,T;W_{\diamond}^{-1,v(p)}(\Omega)) \cap L^{\frac{q}{2}}(0,T;W^{1,v(p)}(\Omega)) \hookrightarrow C([0,T];L^p(\Omega)),$
 (b) $W_0^{1,\frac{q}{2}}(0,T;W_{\diamond}^{-1,v(p)}(\Omega)) \cap L^{\frac{q}{2}}(0,T;W^{1,v(p)}(\Omega)) \hookrightarrow C([0,T];L^p(\Omega)).$

Proof. The embeddings [item 1a](#) and [item 2a](#) follow directly from [Lemma 37](#). The embeddings [item 1b](#) and [item 2b](#) are the restriction of the embeddings from [item 1a](#) and [item 2a](#), respectively, to the subspace $\{\psi : \psi(0) = 0\}$. \square

- Amann, H. (2005). Nonautonomous parabolic equations involving measures. *Journal of Mathematical Sciences*, 130, 4780–8002.
- Bartels, S., & Roubíček, T. (2008). Thermoviscoplasticity at small strains. *ZAMM. Zeitschrift für Angewandte Mathematik und Mechanik. Journal of Applied Mathematics and Mechanics*, 88, 735–754. doi:[10.1002/zamm.200800042](#).
- Bartels, S., & Roubíček, T. (2011). Thermo-visco-elasticity with rate-independent plasticity in isotropic materials undergoing thermal expansion. *ESAIM. Mathematical Modelling and Numerical Analysis*, 45, 477–504. doi:[10.1051/m2an/2010063](#).
- Chelmiński, K., & Racke, R. (2006). Mathematical analysis of thermoplasticity with linear kinematic hardening. *Journal of Applied Analysis*, 12, 37–57. doi:[10.1515/JAA.2006.37](#).
- Dore, G. (1993). L^p regularity for abstract differential equations. In *Functional Analysis and Related Topics, 1991* (pp. 25–38). Springer.
- Gajewski, H., Gröger, K., & Zacharias, K. (1974). *Nichtlineare Operatorgleichungen und Operatordifferentialgleichungen*. Berlin: Akademie-Verlag.
- Grisvard, P. (1985). *Elliptic Problems in Nonsmooth Domains*. Boston: Pitman.
- Gröger, K. (1989). A $W^{1,p}$ -estimate for solutions to mixed boundary value problems for second order elliptic differential equations. *Mathematische Annalen*, 283, 679–687. doi:[10.1007/BF01442860](#).
- Haller-Dintelmann, R., Meyer, C., Rehberg, J., & Schiela, A. (2009). Hölder continuity and optimal control for nonsmooth elliptic problems. *Applied Mathematics and Optimization*, 60, 397–428. doi:[10.1007/s00245-009-9077-x](#).
- Han, W., & Reddy, B. D. (1999). *Plasticity*. New York: Springer.
- Herzog, R., Meyer, C., & Wachsmuth, G. (2011). Integrability of displacement and stresses in linear and nonlinear elasticity with mixed boundary conditions. *Journal of Mathematical Analysis and Applications*, 382, 802–813. doi:[10.1016/j.jmaa.2011.04.074](#).
- Hömberg, D., Meyer, C., Rehberg, J., & Ring, W. (2009/10). Optimal control for the thermistor problem. *SIAM Journal on Control and Optimization*, 48, 3449–3481. doi:[10.1137/080736259](#).

- Ottosen, N., & Ristinmaa, M. (2005). *The Mechanics of Constitutive Modeling*. Elsevier.
- Paoli, L., & Petrov, A. (2012). Global existence result for thermoviscoelastic problems with hysteresis. *Nonlinear Analysis. Real World Applications. An International Multidisciplinary Journal*, 13, 524–542. doi:[10.1016/j.nonrwa.2011.07.018](https://doi.org/10.1016/j.nonrwa.2011.07.018).
- Showalter, R. E. (1997). *Monotone operators in Banach space and nonlinear partial differential equations* volume 49 of *Mathematical Surveys and Monographs*. Providence, RI: American Mathematical Society.
- Simon, J. (1986). Compact sets in the space $L^p(0, T; B)$. *Annali di Matematica Pura ed Applicata*, 146, 65–96. doi:[10.1007/BF01762360](https://doi.org/10.1007/BF01762360).