



RUHR

ECONOMIC PAPERS

Jörg Langbein

Firewood, Smoke and Respiratory Diseases in Developing Countries - the Neglected Role of Outdoor Cooking

Imprint

Ruhr Economic Papers

Published by

Ruhr-Universität Bochum (RUB), Department of Economics
Universitätsstr. 150, 44801 Bochum, Germany

Technische Universität Dortmund, Department of Economic and Social Sciences
Vogelpothsweg 87, 44227 Dortmund, Germany

Universität Duisburg-Essen, Department of Economics
Universitätsstr. 12, 45117 Essen, Germany

RWI Leibniz-Institut für Wirtschaftsforschung
Hohenzollernstr. 1-3, 45128 Essen, Germany

Editors

Prof. Dr. Thomas K. Bauer
RUB, Department of Economics, Empirical Economics
Phone: +49 (0) 234/3 22 83 41, e-mail: thomas.bauer@rub.de

Prof. Dr. Wolfgang Leininger
Technische Universität Dortmund, Department of Economic and Social Sciences
Economics – Microeconomics
Phone: +49 (0) 231/7 55-3297, e-mail: W.Leininger@tu-dortmund.de

Prof. Dr. Volker Clausen
University of Duisburg-Essen, Department of Economics
International Economics
Phone: +49 (0) 201/1 83-3655, e-mail: vclausen@vwl.uni-due.de

Prof. Dr. Roland Döhrn, Prof. Dr. Manuel Frondel, Prof. Dr. Jochen Kluve
RWI, Phone: +49 (0) 201/81 49-213, e-mail: presse@rwi-essen.de

Editorial Office

Sabine Weiler
RWI, Phone: +49 (0) 201/81 49-213, e-mail: sabine.weiler@rwi-essen.de

Ruhr Economic Papers #697

Responsible Editor: Manuel Frondel

All rights reserved. Bochum, Dortmund, Duisburg, Essen, Germany, 2017

ISSN 1864-4872 (online) – ISBN 978-3-86788-811-0

The working papers published in the Series constitute work in progress circulated to stimulate discussion and critical comments. Views expressed represent exclusively the authors' own opinions and do not necessarily reflect those of the editors.

Ruhr Economic Papers #697

Jörg Langbein

**Firewood, Smoke and Respiratory Diseases
in Developing Countries - the Neglected
Role of Outdoor Cooking**

Bibliografische Informationen der Deutschen Nationalbibliothek

Die Deutsche Bibliothek verzeichnet diese Publikation in der deutschen Nationalbibliografie; detaillierte bibliografische Daten sind im Internet über:
<http://dnb.d-nb.de> abrufbar.

Das RWI wird vom Bund und vom Land Nordrhein-Westfalen gefördert.

<http://dx.doi.org/10.4419/86788811>

ISSN 1864-4872 (online)

ISBN 978-3-86788-811-0

Jörg Langbein¹

Firewood, Smoke and Respiratory Diseases in Developing Countries - the Neglected Role of Outdoor Cooking

Abstract

Smoke from cooking in the kitchen is one of the world's leading causes of premature child death, claiming the lives of 500,000 children under five annually. This study analyses the role of outdoor cooking and the prevalence of respiratory diseases among children under five years by means of probit regressions using information from 41 surveys conducted in 30 developing countries from Asia, Africa and Latin America. I find that outdoor cooking reduces respiratory diseases among young children aged 0-4 by around 9 percent, an effect that reaches 13 percent among children aged 0-1. The results suggest that simple behavioral interventions, such as promoting outdoor cooking, can have a substantial impact on health hazards.

JEL Classification: Q53, I12, O13

Keywords: Air pollution; health behavior; energy access

May 2017

¹ Jörg Langbein, RWI. - I am grateful for valuable comments and suggestions by Gunther Bensch, Manuel Frondel, Maike Hohberg, Luciane Lenz, Jörg Peters and Colin Vance. - All correspondence to: Jörg Langbein, RWI, Hohenzollernstr. 1-3, 45128 Essen, Germany, e-mail: joerg.langbein@rwi-essen.de

1 Introduction

About 3 billion people in developing countries rely on firewood or charcoal for their daily cooking purposes (WHO, 2014a). According to the World Health Organization (WHO), smoke-induced diseases are responsible for the death of 4.3 million people every year - more deaths than caused by malaria or tuberculosis - making it one of the most lethal environmental health risks worldwide (WHO, 2014b; Martin, 2011). The largest burden of mortality is borne by women and young children. Among the 4.3 million who die from the consequences of smoke emission each year, 500,000 are children under five that die due to acute respiratory infections (ARI).

Young children are particularly vulnerable for two reasons: First, they are usually with their mothers during the cooking process and thus inhale large loads of particulate emission. In a recent systematic review, it was found that childrens' particulate emission exposure is similar to their mothers' (Balakrishnan et al., 2014). Second, in comparison to adults, the still growing bodies of young children are more susceptible to ARI, leading to a high death rate in this age group (Barnes, 2014).

One proposed remedy are clean cookstoves, such as LPG stoves. However, these cookstoves are expensive and not widespread. Even their more affordable cousin, improved cooking stoves (ICS), which tend to have fewer health and environmental benefits, are only used by around one third of the 2.85 billion people who rely on solid fuel (ESMAP, 2015). In response, many have advocated behavioral change interventions targeting cooking behavior as low-cost interventions to reduce the detrimental health effects from emitted smoke exposure (IOB, 2013; Barnes, 2015; Goodwin, 2015). Such interventions include health education and knowledge dissemination, cooking appliance maintenance training or information on ventilation practices. Yet another behavioral

change is the setting of the cooking location, which is the focus of this paper. Cooking outdoors instead of indoors may have important implications for ventilation and thus smoke exposure (Bensch and Peters, 2015; Dasgupta et al., 2006; Yu, 2011), but has so far been widely neglected in the debate. A review of studies published from 1983 to 2013 suggests that behavioral changes alone may reduce indoor air pollution exposure by 31 to 85 percent (Barnes, 2014). The largest reduction of indoor air pollution exposure was found in a rigorous field study in China, where health education, behavioral activities, and a cookstove intervention were combined (Jin et al., 2006).

This paper explores the correlation of outdoor cooking with acute respiratory infections of children under 5 who live in rural areas of 30 developing countries spanning Asia, Africa and Latin America. So far, only a few articles have explicitly addressed the impact of the cooking location on particulate matter exposure (Nasir et al., 2013; Dasgupta et al., 2009) and respiratory diseases (Albalak et al., 1999; Owusu Boadi and Kuitunen, 2006; Rehfuess et al., 2009; Buchner and Rehfuess, 2015). All of these studies cover only one geographical area of a country and/or use a small dataset. Rehfuess et al. (2009) and Buchner and Rehfuess (2015) are notable exceptions, reporting the effect of outdoor cooking on health for 16 and 9 Sub-Saharan countries.

The source of the present data are the Demographic Health Surveys (DHS), regular cross-sectional surveys in developing countries with a focus on health issues. This study examines 41 surveys conducted in 30 countries, yielding 219,776 observations from the years 2005 to 2014. The decision to cook outdoors is likely jointly determined by other factors. Examples include but are not limited to education, income, cultural factors or climate. This gives rise to potential endogeneity issues, which precludes ascribing a causal interpretation to the effect of outdoor cooking. In order to reduce the omitted

variable bias arising from endogeneity, first a probit model is estimated that controls for confounding variables. In a second step, the probit estimations are subject to several robustness checks, including a matching approach and controlling for smoking behavior. Results are presented on an aggregated level over all countries as well as on a single country level, since cultural differences are one potential source of bias. In order to account for younger children being more susceptible to ARI than older ones, I estimate in a separate model the effect of outdoor cooking on health for children aged 0-1.

It is found that outdoor cooking is associated with a decrease of 0.5 percentage points in the ARI occurrence of children aged 0-4 and with a decrease of 1 percentage point for children aged 0-1 compared to indoor cooking. These coefficients correspond to a reduction due to outdoor cooking of 9 percent of all ARI occurrences for children aged 0-4 and to 13 percent for children aged 0-1. The relationship is significant and remains stable when including control variables or conducting several robustness checks.

The paper is structured as follows: Section 2 describes the literature and some theoretical predictions derived from it. Section 3 provides information on the dataset. The estimation strategy is discussed in Section 4 and results are presented in Section 5. Section 6 concludes.

2 Literature and theoretical prediction

The negative impact of air pollution on health has increased over the last 20 years (Lozano et al., 2012), as has the documentation of it. Combusting solid fuel in-house affects health in various ways and may contribute to acute respiratory infections, stunted growth in children, pneumonia, chronic bronchitis in women, chronic obstructive pulmonary disease

(COPD), cataracts and other visual impairments, cardiovascular diseases, lung cancer, tuberculosis, and perinatal diseases (Po et al., 2011; Ezzati and Kammen, 2002; Amegah et al., 2014; Dherani et al., 2008; McCracken et al., 2012; Hosgood et al., 2010; Bruce et al., 2013). Estimates from the WHO (2014) suggest that the exposure to household air pollution from cooking with solid fuels causes 4.3 million premature deaths annually (WHO, 2014b). The potential harm does not stop once the smoke leaves the house, leading to another 0.5 million premature deaths in 2010 due to ambient air pollution exposure (Brauer et al., 2012; Anenberg et al., 2013; Rao et al., 2012). Several studies highlight that a child’s body is more susceptible to ARI due to still developing lungs and immune systems, which is exacerbated by higher respiration rates among children that lead to around 50 percent more of the polluted air breathed in compared to adults (Pande, 1998; Smith, 2000; Moya et al., 2004).

Key foci of the literature on cookstoves have been on ventilation practices as well as the effect of ambient air pollution on indoor air pollution (Dasgupta et al., 2015, 2009; Ruth et al., 2013). In that context, regular ventilation is generally thought to decrease the particulate matter concentration in the kitchen and other rooms in the house. The same rationale goes for outdoor cooking, where particulate matter directly dissipates in the surroundings. Other studies explicitly analyse the particulate matter concentration for 24 hours on a small scale for indoors and outdoors. Although those studies report a large drop in the concentration level by cooking outside, they note that the level found is still larger than the recommended level by the WHO (Balakrishnan et al., 2002; Rosa et al., 2014). Among the few papers that have explicitly focused on outdoor cooking and health outcomes, Albalak et al. (1999) compare two villages in Bolivia, one cooking outdoors and one indoors. They find that particulate matter exposure is twice as high in

households in the indoor cooking village compared to the outdoor cooking village, translating to a bronchitis reduction of 60 percent. Similar results are found by Owusu and Kuitunen (2006) for Accra, who highlight a stark correlation between children's presence in the indoor cooking area and respiratory diseases (Owusu Boadi and Kuitunen, 2006). Rehfuess et al. (2009) conclude in a cross country study among 16 African countries that the impact of cooking with solids fuels on acute lower respiratory infections is different among children with regards to ventilation practices and the cooking location. Higher ventilation and outdoor cooking decreases the risk of ARI. Buchner and Rehfuess (2015) obtain similar results when analyzing the cooking location in 9 Sub-Saharan African countries. Both studies report average impacts over all countries.

This paper adds to the existing literature by analyzing the issue using a large sample from different countries, thus, yielding broad geographical coverage. The focus is on the link between the cooking location and incidence of infection, controlling for potential confounding factors.

3 Data

I use data from the Demographic Health Surveys (DHS) to assess the relationship between outdoor cooking and health. The DHS are nationally representative, cross-sectional household surveys with a main survey that focuses on general health conditions and demographics and complementary surveys focusing on explicit health topics, such as malaria and HIV.

All countries included in this analysis are low-income or lower middle-income countries in Africa, Latin America, and South-East Asia according to the World Bank classifica-

tions as of October 31, 2016. Due to data regulations, lack of data collection and lack of variation in outdoor cooking, only 30 of 60 countries fit this classification and are included in the analysis, of which 27 are in Africa, one in Asia and two in Latin America. Necessary information on the cooking locations is found in wave 5, wave 6 and wave 7. The analysis is restricted to the main DHS and does not include any other specifically framed questionnaires. Included countries had to meet a threshold of outdoor cooking of at least 10 percent.

Assuming that ambient air pollution is more prevalent in urban regions, I focus on rural regions to reduce this bias. Another reason for the focus on rural areas is that households should be able to decide between indoor and outdoor cooking, whereby available space may be limited in urban areas. Hence, the results only apply to households living in a rural environment and cannot be directly transferred to urban environments.

The main DHS questionnaire asks questions on stove usage, cooking fuels and cooking location but due to a massive lack of responses, the answers on stove usage could not be included in this analysis. Regarding the cooking location, households were asked whether they usually cook in the house, in a separate building, or outside. Multiple answers were not possible. In this analysis, cooking in a separate building and cooking in the house are combined and analyzed as indoor cooking, but an additional robustness check excludes cooking in a separate building, thereby focusing only on the comparison of cooking in the house and outdoors. Cooking fuels are controlled for throughout this analysis. As another robustness test on the effect of outdoor cooking, I restrict the samples to those households using firewood as cooking fuel. This should yield a more precise estimate of the impact of cooking outdoors using solid fuels because cooking with electricity and LPG, which are clean cooking fuels, is mostly done indoors.

Information on the child’s health, answered by the head of household’s wife is merged with the household dataset. The main outcome variable “ARI” is binary and constructed using information on coughing behavior of children aged 0-4 years in the household in the last two weeks before the survey was administered. Mothers were asked to report if the child aged 0-4 years had a cough in the last two weeks before the survey was administered. If this was true, the mother was asked if the coughing was accompanied with fast and heavy breathing due to a problem in the chest and/or nose. When these conditions were met, the pattern was classified as ARI by the DHS program (DHS, 2006).

To obtain results for this age group only, and since there is no data provided for other household members, all households without children age 0-4 years are dropped. The unit of observation are children under the age of 5 and households with several children under 5 appear multiple times in the data.

Overall, 219,776 observations from 30 countries and the years 2005 to 2014 are included. Table A.1 in the supplementary appendix shows the numbers of observations by country and survey year in rural areas.

4 Empirical strategy

Deriving a strictly causal interpretation of the relationship between the cooking location and potential impacts on children’s health in this setup is complicated by the fact that the cooking location is endogenously determined. For example, it may be that better educated household heads or the persons cooking know about the negative health effects of household air pollution and thus cook outdoors. This decision may also be correlated with the household’s earnings. Although I control for education and wealth indicators

as proxies for income and similar confounders, it is not possible to rule out the existence of unobserved variables, such as astuteness or health knowledge, that bias the estimates through their correlation with the outdoor cooking decision.

To estimate the effect of outdoor cooking on the probability that a child suffers from ARI in the last two weeks before the survey, I specify a probit model, in which the binary variable ARI is regressed on the cooking location and a set of control variables.

All waves are pooled and country dummies are included in every estimation to account for country differences. A dummy for the year of the interview is also included to control for time trends. Recognizing that previous research has identified the season as an important driver of ARI, a dummy variable controlling for the occurrence of a rainy season during the time of the interview is constructed (Smith et al., 2011; Paynter et al., 2013; Haspil-Corgan et al., 2004).¹ The variable also takes regional variations of the rainy seasons within the same country into account. However, the interpretation of this variable has to be treated with caution for two reasons: First, it measures whether the interview was conducted during a rainy season but the indicator looked at is retrospective and asks for the occurrence of ARI in the last two weeks before the survey. Second, the survey implementation is not independent of the rainy season and in many cases surveys were only conducted in dry season. Thus, for around half of the countries, there are no observation in both, rainy and dry season and thus, the rainy season dummy is picking up some idiosyncratic country effect. At the household level, control variables for several socio-economic household characteristics, such as ownership of motorbikes, bicycles, cars and/or television, housing wall quality as well as the number of household members, and availability of piped water are added. An additional binary variable is

¹ This was done using information on the rainy seasons from the CIA Factbook. If the needed information could not be found there, other sources were used to obtain information on the seasons.

constructed controlling for regular smoking within the house. The variable takes the value 1 if household members smoke daily or weekly in the house and 0 otherwise. Yet as this variable suffers from a large share of missing observations, this is only included in a robustness check. Regarding the household composition, the age of the household head, the sex of the household head and the education level of the wife are included as controls. At the child level, information on the child’s sex, age and whether the child has ever been vaccinated is included. As with regular smoking, the vaccination indicator suffers from a large number of missing observations and can thus only be included in a robustness check and not in the main specification. Table A.2 in the supplementary appendix for a summary of the variables and the level they are collected. Standard errors are clustered at the household level since children are nested within households.

5 Results

5.1 Descriptive Results

The number of households cooking outdoors varies quite substantially across continents: In Africa, every third household cooks outside, compared to only 15 percent in Asia and 43 percent in Latin America. Due to the dominance of African countries in the data, every third household in the sample cooks outdoors. There are large disparities within Africa, with only 18 percent of the households cooking outdoors in East Africa compared to 43 percent in West Africa (see Table A.3 in the supplementary appendix). There are minor differences in outdoor cooking across seasons. If the interview was conducted in a dry season 38 percent of the households cooked outside, whereas this share amounted

to 31 percent of the households in rainy seasons. Note that the rainy season variable describes whether the region where the household was interviewed in this month was in a rainy season month. Hence, the variable is collected on a regional level, rendering any meaningful interactions with household variables impossible. All following models are estimated with and without the rainy season dummy to check their robustness. The coefficient size and the significance levels remained the same.

Among children aged 0-4 (0-1) around 7 (8) percent suffered from acute respiratory infections in the two weeks preceding the survey. The share is different between the three continents: In African countries, on average 6 percent of the children aged 0-4 suffered from ARI indications, compared to 9 percent in the Philippines as Asian country and 19 percent in the Latin American countries. This share is slightly higher among the subgroup of children aged between 0 and 1, where 8, 9 and 20 percent of the children suffered from ARI indications in Africa, Asia and Latin America, respectively. In West Africa 4 percent (5 percent) of the children aged 0-4 (0-1) suffered from ARI compared to 10 percent (12 percent) in East Africa (all values Table A.3 in the supplementary appendix).

With regards to the control variables, it can be seen that around 92 percent of the included households use firewood as their main cooking fuel and around 16 percent of the wives in the household have finished at least primary education. The average age of the head of the household is around 41 years, while the average household size comprises 7 members. Only 17 percent are female headed households. Daily or weekly smoking is common in 21 percent of the households. Children's gender is split equally between boys and girls, 75 percent had at least one vaccination and 41 percent are aged 0-1 years and 49 percent aged 2-4 years (for all values see Table A.4 in the supplementary appendix).

5.2 Estimation Results

Column 2 of Table 1 presents the marginal effects from a probit model without household characteristics but including dummies for the country, the year of data collection and if the interview was conducted during the rainy season as controls. The model demonstrates that households cooking outdoors had lower occurrence of children’s acute respiratory infections in the two weeks preceding the survey. With a highly significant estimate, cooking outdoors is associated with a decrease in ARI by 0.5 percentage points compared to cooking indoors. This corresponds to a difference of 9 percent in the occurrence of ARI of households that cook outdoors compared to households that cook indoors (Table 1, column 2). This difference increases when restricting the observations to children aged from 0-1 years. In this case I find a reduction of 0.9 percentage points, which is again significant at a 1 percent level in the probit model and corresponds to a difference of 11 percent in the occurrence of ARI (column 3 of Table 1). Further controlling for several socio-economic household characteristics, children characteristics and household fuels in columns 3 and 5 (Table 1) shows that the results are very stable and even increase slightly for children aged 0-1 to 1 percentage point.

Electricity, gas, and kerosene turn out to have a negative association with the occurrence of ARI in the last 2 weeks preceding the survey. This effect is as high as 2.1 percentage points for electricity or 2.2 percentage points for kerosene. There is no difference in ARI occurrence between young children exposed to coal and those exposed to firewood as a fuel source (see Table 1, column 3). The coefficients are very high but as only few households are cooking with kerosene, electricity or gas the point estimates should be treated with caution.

Table 1: ARI occurrence in rural areas (probit results) in the last two weeks preceding the survey^{a,b}

	ARI	ARI	ARI	ARI
	Children	Children	Children	Children
	0-4 years	0-4 years	0-1 years	0-1 years
	<i>margins</i>	<i>margins</i>	<i>margins</i>	<i>margins</i>
Outdoor cooking	-0.005*** (0.00)	-0.005*** (0.00)	-0.009*** (0.00)	-0.010*** (0.00)
Electricity as cooking fuel		-0.022*** (0.01)		-0.045*** (0.01)
Gas as cooking fuel		-0.014*** (0.00)		-0.032*** (0.01)
Kerosene as cooking fuel		-0.029*** (0.01)		-0.033*** (0.01)
Coal, Lignite / Charcoal as cooking fuel		-0.000 (0.00)		-0.003 (0.00)
Other cooking fuel		0.007 (0.00)		0.003 (0.00)
Child is female		-0.003*** (0.00)		-0.006*** (0.00)
Observations	219,776	218,970	90,747	90,416
Country dummies	Yes	Yes	Yes	Yes
Year of data collection dummies	Yes	Yes	Yes	Yes
Interview in rainy season dummy	Yes	Yes	Yes	Yes
Household characteristics	No	Yes	No	Yes


Note: *, **, *** indicate p-values at the 10 percent level, 5 percent level and 1 percent level, respectively. All estimations are clustered on household level and standard errors are in brackets. Table A.5 in the supplementary appendix gives the coefficients to the marginal effects. Using wood as cooking fuel is the baseline category for the variable on cooking fuels and outdoor cooking is analysed in respect to indoor cooking.

Source: DHS all country dataset from 2005-2014.

Figure 1 presents estimates of the mean effect of outdoor cooking on ARI for children aged 0-1 based on models estimated for all countries separately.² In this case, 20 countries exhibit a negative correlation between outdoor cooking and ARI. Of these, 10 of the estimated effects are statistically significant. There is no country with a significantly positive relationship between ARI and outdoor cooking.

² Similar results are obtained for the children aged 0-4 years.

Figure 1: Effect size of outdoor cooking on ARI for children aged 0-1 (Probit estimation with controls)



Note: Bars signal 95 percent confidence intervals. Standard errors are clustered on household level. Congo refers to the Republic of Congo and Congo (Dem. Rep) refers to the Democratic Republic of the Congo.

Source: DHS all country dataset from 2005-2014.

Although the general trend shows a negative relationship between outdoor cooking and ARI, the relationship on the country level is heterogeneous. It is not entirely clear why this is the case, but the following provides some reasoning for the result. In some countries, many households have fixed built-in versions of cookstoves. In most of the Sub-Saharan African countries, portable stoves that can be used for cooking both inside and outside are prominent but some countries like Ethiopia and Uganda have a long tradition of using built-in stoves. With built-in stoves, households cook indoors but the emission in the cooking process is reduced. Thus, the particulate matter concentration may be less indoors than cooking outdoors without a stove (ESMAP, Energy Sector Management Assistance Program, 2015). Built-in stoves are also popular in Honduras, where clean cooking solutions for stationary stoves have received large political support (ESMAP, Energy Sector Management Assistance Program, 2015).

In a next step, the robustness of the results is assessed by limiting the dataset to firewood-using households only, and excluding households that cook in a separate building (see Table 2 and Tables A.6 and A.7 in the supplementary appendix).

Table 2: ARI in the last 2 weeks preceding the survey in rural areas limited to firewood cooking households and excluding separate building cooking households ^{a, b}

	ARI	ARI	ARI	ARI
	Children	Children	Children	Children
	0-4 years	0-1 years	0-4 years	0-1 years
	<i>margins</i>	<i>margins</i>	<i>margins</i>	<i>margins</i>
	Firewood using households only	Firewood using households only	Seperate building households excluded	Seperate building households excluded
Outdoor cooking	-0.006*** (0.00)	-0.011*** (0.00)	-0.009*** (0.00)	-0.013*** (0.00)
Observations	203,068	83,440	135,542	56,141
Country dummies	Yes	Yes	Yes	Yes
Year of data collection dummies	Yes	Yes	Yes	Yes
Interview in rainy season dummy	Yes	Yes	Yes	Yes
Household characteristics	Yes	Yes	Yes	Yes

Note: *, **, *** indicate p-values of a 10 percent level, 5 percent level and 1 percent level, respectively. All estimations are clustered on household level and standard errors are in brackets. Tables A.6 and A.7 in the supplementary appendix give the coefficients to the marginal effects.

Source: DHS all country dataset from 2005-2014.

Dropping all households that do not use firewood as the main cooking fuel reduces the sample to 203,068 households. Estimating a probit model yields a larger negative coefficient, providing further evidence for the negative relationship between outdoor cooking and ARI among children (Table 2). Table A.6 in the supplementary appendix shows the full model. For households using firewood as the main cooking fuel, the negative association increases from -0.5 percentage points to -0.6 percentage points in the model with all controls (Table 2, column 1). Among children aged 0-1 the model shows an increase in the outdoor cooking coefficient from -1 percentage point to -1.1 percentage point. All relevant estimates remained statistically significant on a 1 percent level. The two principal models are re-estimated with only the households that cook in the house or outdoors, thereby excluding the households that cook in a separate building. The respective coefficients increase in size compared to the same estimations that included households cooking in separate building. Children aged 0-4 (0-1) living in households that cook outdoors suffered a -0.9 (-1.3) percentage point lower ARI incidence than those who live in households that cook indoors in the main house (see Table 2, column 3 and 4). All models are further estimated using a matching approach with nearest neighbour

matching without replacement. The marginal effects and significance levels are similar to those obtained using probit methods (see Table A.8 in the supplementary appendix for the full model). As a last robustness check, I estimate the principal models with the frequency of smoking in the household and vaccination of children as control variables. However, as both variables have large number of missing observations, it is not possible to include the two of them in the main specifications. In fact, it is even not possible to have both as control variables in the same regression as this would result in the dropping of around 75 percent of the observations. Table A.9 in the supplementary appendix reports the results on including frequency of smoking in the household as well as vaccination status of children separately. With the caveat of a largely reduced sample in mind, the results including vaccination as control variable are a bit lower but in general very similar to the main results. Those including frequency of smoking in the households as additional control yield an increased negative coefficient of -1.6 percentage points for children aged 0-1. For this sample, a reduction of -1.6 percentage points is corresponding to a decrease of 27 percent of ARI among children aged 0-1.

6 Conclusion

This paper analyzes the relationship between outdoor cooking and children's acute respiratory infections in 30 developing countries over 3 continents. The results demonstrate a large and statistically significant relationship between outdoor cooking and the occurrence of ARI with an emphasis on children aged 0-1.

Although it was not possible to definitely isolate a causal relationship, the results point to the fact that outdoor cooking has a substantial association with children's health. It

is calculated that the effect size is as large as 9 percent for children between 0 and 4 when controlling for potential confounders. The effect size is even larger for children aged between 0 and 1, reaching a magnitude of 13 percent. This larger effect size for the very young children is attributed to the fact that the childrens' body is more susceptible to particulate smoke emission at this age and children are more likely to be with their mothers. Estimating the main specifications excluding those households that cook in a separate building yields a larger coefficient on outdoor cooking compared to in-house cooking only. A last robustness check, including smoking frequency in the households, shows that the effect size increases starkly to -1.6 percentage points when controlling for this confounder for the children aged 0-1.

The overall results are complementary to those found by Buchner and Rehfuess (2015) and Rehfuess et al. (2009), as both find a significant negative association between outdoor cooking and ARI. This paper extends the analysis not only by including more countries but also by exploiting a large degree of heterogeneity between countries with respect to the incidence of cooking outdoors and the associated impact on children's health.

Several caveats of the analysis point to areas where future research is needed: The dataset does not contain information on the type of cooking stoves used, fuel stacking, or other potential cofounders, making it impossible to assess the impact of those factors on ARI. The analysis thus rests on the assumption that any omitted variable is not systematic and does not bias the results extensively. I also have no information on the children's presence at the cooking site. It is likely that the direct effect on childrens' health is higher if all children remain in close proximity to the cooking site as suggested by studies that only examine the particulate emission (Rosa et al. (2014), for example). This may partly explain the difference of the magnitude in the effects for children aged 0-1

and children aged 0-4. However, this approach has the virtue of exploiting the occurrence of ARI, compared to other studies that examine the emittance of particulate emission only.

In this respect the results can be related to the influence of ARI on child mortality. Smith et al. (2014) report that around 466,000 children died in 2010 due to respiratory infections. Assuming that decreases in deaths from ARI are proportional to decreases in its incidence, the estimated 9 percent reduction in ARI from cooking outdoors implies that 41,900 fewer children would die annually if everyone cooked outdoors.

The results of this study fit in the general literature on health hazards from particulate emission by significantly contributing to the scarce evidence of the impact of outdoor cooking on health hazards. Several policy implications emerge. Most importantly, the results demonstrate that simple behavioral interventions, such as promoting outdoor cooking, can have a substantial impact on health hazards. Of course, not everyone can cook outside the whole year due to unfavorable seasonal conditions. Such constraints should be taken into account when formulating policy, targeting behavioral interventions in those regions where there is scope for moving the cooking location outdoors. With regards to child protection, raising awareness of the detrimental effects of smoke emission on children and the benefits of leaving the children only a few meters away from the cooking place is important, particularly in countries where young children are always with their mothers. Behavioral interventions are nevertheless no panacea and should be complemented by the distribution of clean cookstoves. Yet, adopting cleaner fuels, such as LPG, has proven to be difficult and will be a long-term goal. In the meantime, policy effort should concentrate on promoting behavioral strategies and ventilation, with the extreme case of outdoor cooking being an important component.

In this regard, more solid evidence on the influence of different behavioral factors such as ventilation practices, outdoor cooking or general cooking behavior on health issues in general and children's health in particular is needed. The combination of strategies targeting cookstove adoption and low-cost health knowledge interventions is a promising path for policy makers. Applying state of the art techniques, such as randomized controlled trials, to evaluate such interventions would be beneficial.

References

- Albalak, R., Frisancho, A. and Keeler, G. (1999). Domestic biomass fuel combustion and chronic bronchitis in two rural Bolivian villages, *Thorax* **54**(11): 1004–1008.
- Amegah, A. K., Quansah, R. and Jaakkola, J. J. K. (2014). Household air pollution from solid fuel use and risk of adverse pregnancy outcomes: a systematic review and meta-analysis of the empirical evidence, *PloS one* **9**(12): e113920.
- Anenberg, S. C., Balakrishnan, K., Jetter, J., Masera, O., Mehta, S., Moss, J. and Ramanathan, V. (2013). Cleaner cooking solutions to achieve health, climate, and economic cobenefits, *Environmental Science & Technology* **47**(9): 3944–3952.
- Balakrishnan, K., Mehta, S., Ghosh, S., Johnson, M., Brauer, M., Zhang, J., Naeher, L. and Smith, K. R. (2014). Review 5: Population levels of household air pollution and exposures, in WHO, World Health Organization (ed.), *Indoor air quality guidelines: household fuel combustion*, Geneva, Switzerland. URL: <http://www.who.int/indoorair/publications/household-fuel-combustion/en/> [last accessed on 23/01/2016].
- Balakrishnan, K., Sankar, S., Parikh, J., Padmavathi, R., Srividya, K., Venugopal, V., Prasad, S. and Pandey, V. L. (2002). Daily average exposures to respirable particulate matter from combustion of biomass fuels in rural households of southern India, *Environmental Health Perspectives* **110**(11): 1069–1075.
- Barnes, B. R. (2014). Behavioural change, indoor air pollution and child respiratory health in developing countries: a review, *International Journal of Environmental Research and Public Health* **11**(5): 4607–4618.
- Barnes, B., Rosenbaum, J., Mehta, S., Williams, K. N., Jagoe, K. and Graham, J. (2015). Behavior change communication: a key ingredient for advancing clean cooking, *Journal of Health Communication* **20**(sup1): 3–5.
- Bensch, G. and Peters, J. (2015). The intensive margin of technology adoption-experimental evidence on improved cooking stoves in rural Senegal, *Journal of Health Economics* **42**: 44–63.
- Brauer, M., Amann, M., Burnett, R. T., Cohen, A., Dentener, F., Ezzati, M., Henderson, S. B., Krzyzanowski, M., Martin, R. V., van Dingenen, R. and et al. (2012). Exposure assessment for estimation of the global burden of disease attributable to outdoor air pollution, *Environmental Science & Technology* **46**(2): 652–660.
- Bruce, N. G., Dherani, M. K., Das, J. K., Balakrishnan, K., Adair-Rohani, H., Bhutta, Z. A. and Pope, D. (2013). Control of household air pollution for child survival: estimates for intervention impacts, *BMC Public Health* **13 Suppl 3**: S8.
- Buchner, H. and Rehfuess, E. A. (2015). Cooking and season as risk factors for acute lower respiratory infections in African children: a cross-sectional multi-country analysis, *PloS one* **10**(6): e0128933.
- Dasgupta, S., Huq, M., Khaliqzaman, M., Pandey, K. and Wheeler, D. (2006). Who suffers from indoor air pollution? Evidence from Bangladesh, *Health Policy and Planning* **21**(6): 444–458.

- Dasgupta, S., Martin, P. and Samad, H. A. (2015). Lessons from rural Madagascar on improving air quality in the kitchen, *The Journal of Environment & Development* **24**(3): 345–369.
- Dasgupta, S., Wheeler, D., Huq, M. and Khaliqzaman, M. (2009). Improving indoor air quality for poor families: a controlled experiment in Bangladesh, *Indoor Air* **19**(1): 22–32.
- Dherani, M., Pope, D., Mascarenhas, M., Smith, K. R., Weber, M. and Bruce, N. (2008). Indoor air pollution from unprocessed solid fuel use and pneumonia risk in children aged under five years: a systematic review and meta-analysis, *Bulletin of the World Health Organization* **86**(5): 390–398.
- DHS, Demographic and Health Survey (2006). *Guide to DHS statistics – Demographic and Health Survey methodology*, United States Agency International Development, Maryland.
- ESMAP, Energy Sector Management Assistance Program (2015). The state of the global clean and improved cooking sector, *Technical Report 007/15*, The International Bank for Reconstruction and Development and the World Bank, Washington, DC.
- Ezzati, M. and Kammen, D. M. (2002). The health impacts of exposure to indoor air pollution from solid fuels in developing countries: knowledge, gaps, and data needs, *Environmental Health Perspectives* **110**(11): 1057–1068.
- Goodwin, N. J., O’Farrell, S. E., Jagoe, K., Rouse, J., Roma, E., Biran, A. and Finkelstein, E. A. (2015). Use of behavior change techniques in clean cooking interventions: a review of the evidence and scorecard of effectiveness, *Journal of Health Communication* **20**(sup1): 43–54.
- Haspil-Corgan, T. A., Tewari, M., Low, R. B. and Gauff, W. R. (2004). The effects of weather on pneumonia incidence, *Annals of Emergency Medicine* **44**(4): S37.
- Hosgood, H. D., Boffetta, P., Greenland, S., Lee, Y.-C. A., McLaughlin, J., Seow, A., Duell, E. J., Andrew, A. S., Zaridze, D., Szeszenia-Dabrowska, N., Rudnai, P. and et al. (2010). In-home coal and wood use and lung cancer risk: a pooled analysis of the International Lung Cancer Consortium, *Environmental Health Perspectives* **118**(12): 1743–1747.
- IOB, Ministry of Foreign Affairs of the Netherlands (2013). *Renewable energy: access and impact – A systematic literature review of the impact on livelihoods of interventions providing access to renewable energy in developing countries*, IOB, Den Haag.
- Jin, Y., Ma, X., Chen, X., Cheng, Y., Baris, E. and Ezzati, M. (2006). Exposure to indoor air pollution from household energy use in rural China: the interactions of technology, behavior, and knowledge in health risk management, *Social Science & Medicine (1982)* **62**(12): 3161–3176.
- Lozano, R., Naghavi, M., Foreman, K., Lim, S., Shibuya, K., Aboyans, V., Abraham, J., Adair, T., Aggarwal, R., Ahn, S. Y., AlMazroa, M. A. and et al. (2012). Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: A systematic analysis for the Global Burden of Disease Study 2010, *The Lancet* **380**(9859): 2095–2128.
- Martin, W. J. n., Glass, R. I., Balbus, J. M. and Collins, F. S. (2011). Public health. A major environmental cause of death, *Science* **334**(6053): 180–181.

- McCracken, J. P., Wellenius, G. A., Bloomfield, G. S., Brook, R. D., Tolunay, H. E., Dockery, D. W., Rabadan-Diehl, C., Checkley, W. and Rajagopalan, S. (2012). Household air pollution from solid fuel use: evidence for links to CVD, *Global Heart* **7**(3): 223–234.
- Moya, J., Bearer, C. F. and Etzel, R. A. (2004). Children’s behavior and physiology and how it affects exposure to environmental contaminants, *Pediatrics* **113**(Supplement 3): 996–1006.
- Nasir, Z. A., Colbeck, I., Ali, Z. and Ahmad, S. (2013). Indoor particulate matter in developing countries: a case study in Pakistan and potential intervention strategies, *Environmental Research Letters* **8**(2): 024002.
- Owusu Boadi, K. and Kuitunen, M. (2006). Factors affecting the choice of cooking fuel, cooking place and respiratory health in the Accra metropolitan area, Ghana, *Journal of Biosocial Science* **38**(3): 403–412.
- Pande, J. N. (1998). *Respiratory medicine in the tropics*, Oxford University Press, Delhi and New York.
- Paynter, S., Weinstein, P., Ware, R. S., Lucero, M. G., Tallo, V., Nohynek, H., Barnett, A. G., Skelly, C., Simoes, E. A. F., Sly, P. D. and Williams, G. (2013). Sunshine, rainfall, humidity and child pneumonia in the tropics: time-series analyses, *Epidemiology and Infection* **141**(6): 1328–1336.
- Po, J. Y. T., FitzGerald, J. M. and Carlsen, C. (2011). Respiratory disease associated with solid biomass fuel exposure in rural women and children: systematic review and meta-analysis, *Thorax* **66**(3): 232–239.
- Rao, S., Chirkov, V., Dentener, F., van Dingenen, R., Pachauri, S., Purohit, P., Amann, M., Heyes, C., Kinney, P., Kolp, P., Klimont, Z. and et al. (2012). Environmental modeling and methods for estimation of the global health impacts of air pollution, *Environmental Modeling & Assessment* **17**(6): 613–622.
- Rehfuess, E. A., Tzala, L., Best, N., Briggs, D. J. and Joffe, M. (2009). Solid fuel use and cooking practices as a major risk factor for ALRI mortality among African children, *Journal of Epidemiology and Community Health* **63**(11): 887–892.
- Rosa, G., Majorin, F., Boisson, S., Barstow, C., Johnson, M., Kirby, M., Ngabo, F., Thomas, E. and Clasen, T. (2014). Assessing the impact of water filters and improved cook stoves on drinking water quality and household air pollution: a randomised controlled trial in Rwanda, *PloS one* **9**(3): e91011.
- Ruth, M., Maggio, J., Whelan, K., DeYoung, M., May, J., Peterson, A. and Paterson, K. (2013). Kitchen 2.0: Design guidance for healthier cooking environment, *International Journal for Service Learning in Engineering* **Special Issue**: 151–169.
- Smith, K. R. (2000). Indoor air pollution in developing countries and acute lower respiratory infections in children, *Thorax* **55**(6): 518–532.
- Smith, K. R., Bruce, N., Balakrishnan, K., Adair-Rohani, H., Balmes, J., Chafe, Z., Dherani, M., Hosgood, H. D., Mehta, S., Pope, D. and Rehfuess, E. (2014). Millions dead: how do we know and what does it mean? Methods used in the comparative risk assessment of household air pollution, *Annual Review of Public Health* **35**: 185–206.

- Smith, K. R., McCracken, J. P., Weber, M. W., Hubbard, A., Jenny, A., Thompson, L. M., Balmes, J., Diaz, A., Arana, B. and Bruce, N. (2011). Effect of reduction in household air pollution on childhood pneumonia in Guatemala (RESPIRE): a randomised controlled trial, *The Lancet* **378**(9804): 1717–1726.
- WHO, World Health Organization (2014a). Burden of disease from household air pollution for 2012. URL: http://www.who.int/phe/health_topics/outdoorair/databases/en/ [last accessed on 11/11/2016].
- WHO, World Health Organization (ed.) (2014b). *Indoor air quality guidelines: household fuel combustion*, Geneva, Switzerland.
- Yu, F. (2011). Indoor air pollution and children’s health: net benefits from stove and behavioral interventions in rural China, *Environmental and Resource Economics* **50**(4): 495–514.

Appendix

Table A.1: Sample description of households in rural areas

Country	Continent(Region)	Survey year(s)	Observations
Benin	Africa(West)	2012	7,450
Burkina Faso	Africa (West)	2010	9,759
Cameroon	Africa (Central/South)	2011	4,454
Comoros	Africa (East)	2012	1,672
Cote d'Ivoire	Africa (West)	2012	3,777
Democratic Republic of the Congo	Africa (Central/South)	2014	9,211
Ethiopia	Africa (East)	2011	7,731
Gabon	Africa (Central/South)	2012	1,485
Gambia	Africa (West)	2013	4,600
Ghana	Africa (West)	2008	4,622
Guinea	Africa (West)	2012	3,841
Haiti	Latin America	2006,2012	7,156
Honduras	Latin America	2012	5,110
Lesotho	Africa (Central/South)	2009,2014	3,848
Liberia	Africa (West)	2013	6,099
Madagascar	Africa (East)	2009	8,672
Malawi	Africa (East)	2010	12,791
Mali	Africa (West)	2013	6,739
Namibia	Africa (Central/South)	2007,2013	4,280
Niger	Africa (West)	2012	8,030
Nigeria	Africa (West)	2006,2013	33,755
Philippines	Asia	2008,2013	5,659
Republic of the Congo	Africa (Central/South)	2012	5,118
Rwanda	Africa (East)	2010	10,480
Senegal	Africa (West)	2011	6,955
Sierra Leone	Africa (West)	2013	8,882
Togo	Africa (West)	2014	3,672
Uganda	Africa (East)	2006,2011	8,457
Zambia	Africa (East)	2007,2014	9,562
Zimbabwe	Africa (Central/South)	2005,2011	5,909
Total			219,776

Source: DHS all country dataset from 2005–2014.

Table A.2: Description of variables included in the analysis

	Analysis level
Acute respiratory infection in the last 2 weeks (constructed from the questions if children had coughing in the last two weeks preceeding the interview in combination with short, rapid breaths and problems in the chest), children gender, children vaccination	Children level
Type of cooking fuel used in the household, household cooking outdoors, age of head of households, head of household is female, number of household members, wife has at least primary education, year of the interview, house build out of high quality material (like stone), household owns a bike, household owns a motorcycle, household owns a car, household has a television, household has access to piped water, frequency of smoking in the household	Household Level
Rainy season	Regional level
Country dummies, interview year dummies	Country level

Table A.3: Outdoor cooking and acute respiratory infections (ARI) prevalence for households in rural areas

Country	Share of households cooking outdoors	Share of ARI prevalence	Share of ARI for children aged 0-1	Number of Observations of children aged 0-4 (of children aged 0-1)
Africa				
Burkina Faso	0.71 (0.45)	0.02 (0.13)	0.02(0.15)	9,759 (3,999)
Benin	0.59 (0.49)	0.01 (0.11)	0.02 (0.12)	7,450 (2,944)
Cameroon	0.24 (0.43)	0.08 (0.27)	0.10 (0.30)	4,454 (1,958)
Comoros	0.22 (0.41)	0.03 (0.18)	0.04 (0.19)	1,672 (690)
Cote d'Ivoire	0.49 (0.50)	0.04 (0.20)	0.05 (0.21)	3,777 (1,628)
Democratic Republic of the Congo	0.34 (0.47)	0.09 (0.29)	0.12 (0.33)	9,211 (3,697)
Ethiopia	0.15 (0.36)	0.08 (0.28)	0.10 (0.31)	7,731 (2,921)
Gabon	0.31 (0.46)	0.13 (0.33)	0.12 (0.32)	1,485 (644)
Ghana	0.46 (0.50)	0.06 (0.23)	0.06 (0.24)	4,622 (1,938)
Gambia	0.11 (0.32)	0.05 (0.21)	0.06 (0.24)	4,600 (2,092)
Guinea	0.34 (0.47)	0.08 (0.26)	0.10 (0.30)	3,841 (1,642)
Liberia	0.58 (0.49)	0.11 (0.32)	0.14 (0.35)	6,099 (2,512)
Lesotho	0.56 (0.50)	0.08 (0.27)	0.09 (0.28)	3,848 (1,721)
Madagascar	0.11 (0.32)	0.03 (0.17)	0.04 (0.20)	8,672 (3,428)
Mali	0.18 (0.39)	0.02 (0.14)	0.02 (0.14)	6,739 (2,714)
Malawi	0.29 (0.46)	0.09 (0.28)	0.10 (0.30)	12,791 (5,122)
Niger	0.81 (0.39)	0.05 (0.22)	0.07 (0.26)	8,030 (3,196)
Nigeria	0.23 (0.42)	0.03 (0.17)	0.04 (0.19)	33,755 (14,287)
Namibia	0.37 (0.48)	0.08 (0.26)	0.08 (0.27)	4,280 (1,903)
Republic of the Congo	0.46 (0.50)	0.06 (0.24)	0.08 (0.26)	5,118 (2,138)
Rwanda	0.19 (0.40)	0.06 (0.23)	0.08 (0.27)	10,480 (3,937)
Sierra Leone	0.58 (0.49)	0.07 (0.26)	0.09 (0.28)	8,882 (3,858)
Senegal	0.11 (0.32)	0.05 (0.22)	0.05 (0.23)	6,955 (2,918)
Togo	0.63 (0.48)	0.05 (0.22)	0.05 (0.22)	3,672 (1,536)
Uganda	0.21 (0.41)	0.23 (0.42)	0.27 (0.45)	8,457 (3,484)
Zambia	0.38 (0.48)	0.05 (0.23)	0.06 (0.25)	9,562 (3,937)
Zimbabwe	0.11 (0.32)	0.07 (0.26)	0.07 (0.26)	5,909 (2,617)
<i>Overall Africa</i>	0.34 (0.47)	0.06 (0.23)	0.08 (0.26)	201,851 (83,461)
Asia				
Philippines	0.15 (0.36)	0.09 (0.28)	0.09 (0.28)	5,659 (2,207)
<i>Overall Asia</i>	0.15 (0.36)	0.09 (0.28)	0.09 (0.28)	5,659 (2,207)
Latin America				
Honduras	0.12 (0.32)	0.18 (0.39)	0.19 (0.39)	5,110 (2,141)
Haiti	0.61 (0.49)	0.20 (0.40)	0.20 (0.40)	7,156 (2,938)
<i>Overall Latin America</i>	0.43 (0.49)	0.19 (0.39)	0.20 (0.40)	12,266 (5,079)
Overall	0.34 (0.47)	0.07 (0.25)	0.08 (0.27)	219,776 (90,747)

Note: Descriptive variables are weighted with official DHS survey weights. Standard deviations in brackets.
Source: DHS all country dataset from 2005–2014.

Table A.4: Descriptive statistics

	Observation	Mean
<i>Household characteristics</i>		
Share of households (in percent) cooking with	213,271	
Electricity	860	0.40
Gas	2,307	1.08
Firewood	196,241	92.01
Charcoal	10,337	4.85
Other	1,561	0.37
Kerosene	1,963	0.92
Wife has at least primary education (=1 if yes)	219,776	0.16 (0.37)
Household has access to piped water	219,643	0.17 (0.37)
House built out of high quality material (=1 if yes)	219,706	0.27 (0.44)
Age of head of household	219,685	40.75 (13.28)
Head of household is female (=1 if yes)	219,776	0.17 (0.37)
Number of household member	219,776	7.12 (4.15)
Household owns bike (=1 if yes)	219,698	0.33 (0.54)
Household owns motorcycle (=1 if yes)	219,678	0.17 (0.46)
Household owns car (=1 if yes)	219,705	0.03 (0.34)
Household owns television (=1 if yes)	219,738	0.14 (0.35)
Regular smoking in the house (=1 if yes)	122,391	0.21 (0.41)
<i>Children characteristics</i>		
Children aged 0-1 years (=1 if yes)	219,776	0.41 (0.49)
Gender of child (=1 if female)	219,776	0.50 (0.50)
Child has ever had a vaccination (=1 if yes)	111,959	0.75 (0.43)

Note: Descriptive variables are weighted with official DHS survey weights. Standard deviations in brackets.

Source: DHS all country dataset from 2005–2014.

Table A.5: Probit estimation of ARI in rural areas with coefficients and marginal effects

	ARI	ARI	ARI	ARI	ARI	ARI	ARI	ARI	ARI
	Children	Children	Children	Children	Children	Children	Children	Children	Children
	0-4 years	0-4 years	0-4 years	0-4 years	0-1 years	0-1 years	0-1 years	0-1 years	0-1 years
	margins	margins	margins	margins	margins	margins	margins	margins	margins
Outdoor cooking	-0.039*** (0.01)	-0.005*** (0.00)	-0.044*** (0.01)	-0.005*** (0.00)	-0.067*** (0.02)	-0.009*** (0.00)	-0.074*** (0.02)	-0.010*** (0.00)	-0.074*** (0.02)
Electricity as cooking fuel			-0.204** (0.08)	-0.022*** (0.01)			-0.420*** (0.13)	-0.045*** (0.01)	-0.420*** (0.13)
Gas as cooking fuel			-0.118*** (0.04)	-0.014*** (0.00)			-0.268*** (0.07)	-0.032*** (0.01)	-0.268*** (0.07)
Kerosene as cooking fuel			-0.279*** (0.07)	-0.029*** (0.01)			-0.274*** (0.10)	-0.033*** (0.01)	-0.274*** (0.10)
Coal, Lignite / Charcoal as cooking fuel			-0.002 (0.02)	-0.000 (0.00)			-0.024 (0.03)	-0.003 (0.00)	-0.024 (0.03)
Other cooking fuel			0.055 (0.06)	0.007 (0.01)			0.022 (0.08)	0.003 (0.01)	0.022 (0.08)
Child is female			-0.026*** (0.01)	-0.003*** (0.00)			-0.040*** (0.01)	-0.006*** (0.00)	-0.040*** (0.01)
Constant	5.660 (5.13)		4.412 (5.17)		-5.569 (6.91)		-8.846 (6.97)		-8.846 (6.97)
Observations	219,776	219,776	218,970	218,970	90,747	90,747	90,416	90,416	90,416
Country dummies	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Year of data collection dummies	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Interview in rainy season dummy	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Household characteristics	No	No	Yes	Yes	No	No	Yes	Yes	Yes

Note: *, **, *** indicate p-values of a 10 percent level, 5 percent level and 1 percent level, respectively. All estimations are clustered on the household level and standard errors are in brackets.

Source: DHS all country dataset from 2005-2014.

Table A.6: Probit estimation of ARI in rural areas with coefficients and marginal effects for firewood using households only

	ARI	ARI	ARI	ARI	ARI	ARI	ARI	ARI
	Children	Children	Children	Children	Children	Children	Children	Children
	0-4 years	0-4 years	0-4 years	0-4 years	0-1 years	0-1 years	0-1 years	0-1 years
		margins		margins		margins		margins
Outdoor cooking	-0.050*** (0.01)	-0.006*** (0.00)	-0.051*** (0.01)	-0.006*** (0.00)	-0.073*** (0.02)	-0.010*** (0.00)	-0.075*** (0.02)	-0.011*** (0.00)
Constant	1.256 (5.39)		.0394 (5.42)		-11.198 (7.22)		-13.319* (7.26)	
Observations	203,068	203,068	202,346	202,346	83,733	83,733	83,440	83,440
Country dummies	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Year of data collection dummies	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Interview in rainy season dummy	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Household characteristics	No	No	Yes	Yes	No	No	Yes	Yes

Note: *, **, *** indicate p-values of a 10 percent level, 5 percent level and 1 percent level, respectively. All estimations are clustered on the household level and standard errors are in brackets.

Source: DHS all country dataset from 2005-2014.

Table A.7: Probit estimation of ARI for households cooking in the main house or outdoors in rural areas with coefficients and marginal effects

	ARI	ARI	ARI	ARI	ARI	ARI	ARI	ARI	ARI
	Children	Children	Children	Children	Children	Children	Children	Children	Children
	0-4 years	0-4 years	0-4 years	0-4 years	0-4 years	0-1 years	0-1 years	0-1 years	0-1 years
	margins	margins	margins	margins	margins	margins	margins	margins	margins
Outdoor cooking	-0.067*** (0.02)	-0.008*** (0.00)	-0.076*** (0.02)	-0.009*** (0.00)	-0.083*** (0.02)	-0.011*** (0.00)	-0.097*** (0.02)	-0.013*** (0.00)	-0.013*** (0.00)
Constant	9.789 (6.49)		8.121 (6.54)		2.055 (8.79)		-2.6888 (8.89)		
Observations	136,098	136,098	135,542	135,542	56,366	56,366	56,141	56,141	56,141
Country dummies	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Year of data collection dummies	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Interview in rainy season dummy	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Household characteristics	No	No	Yes	Yes	No	No	Yes	Yes	Yes

Note: *, **, *** indicate p-values of a 10 percent level, 5 percent level and 1 percent level, respectively. All estimations are clustered on the household level and standard errors are in brackets.
Source: DHS all country dataset from 2005–2014.

Table A.8: Estimation of ARI in rural areas with coefficients and marginal effects (matched results)

	ARI	ARI	ARI	ARI	ARI	ARI	ARI	ARI	ARI
	Children	Children	Children	Children	Children	Children	Children	Children	Children
	0-4 years	0-4 years	0-4 years	0-4 years	0-4 years	0-1 years	0-1 years	0-1 years	0-1 years
	margins	margins	margins	margins	margins	margins	margins	margins	margins
Outdoor cooking	-0.037*** (0.01)	-0.005*** (0.00)	-0.044*** (0.01)	-0.005*** (0.00)	-0.067*** (0.02)	-0.009*** (0.00)	-0.075*** (0.02)	-0.011*** (0.00)	-0.011*** (0.00)
Constant	8.756 (5.93)		2.074 (6.09)		-6.711 (8.13)		-8.958 (8.20)		
Observations	147,637	147,637	147,498	147,498	61,148	61,148	61,090	61,090	61,090
Country dummies	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Year of data collection dummies	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Interview in rainy season dummy	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Household characteristics	No	No	Yes	Yes	No	No	Yes	Yes	Yes

Note: *, **, *** indicate p-values of a 10 percent level, 5 percent level and 1 percent level, respectively. All estimations are clustered on the household level and standard errors are in brackets.

Source: DHS all country dataset from 2005-2014.

Table A.9: Estimation of ARI in rural areas with coefficients and marginal effects with vaccination and frequency of smoking as further control variables

	ARI	ARI	ARI	ARI	ARI	ARI	ARI
	Children	Children	Children	Children	Children	Children	Children
	0-4 years	0-4 years	0-1 years	0-1 years	0-4 years	0-1 years	0-1 years
		margins		margins		margins	margins
Regular smoking in the house							
Outdoor cooking	-0.083*** (0.01)	-0.009*** (0.00)	-0.126*** (0.02)	-0.016*** (0.00)	-0.031** (0.01)	-0.004** (0.00)	-0.051** (0.02)
Observations	122,380	122,380	50,313	50,313	111,945	111,945	38,012
Country dummies	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Year of data collection dummies	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Interview in rainy season dummy	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Household characteristics	No	No	Yes	Yes	No	Yes	Yes
Regular smoking in house	Yes	Yes	Yes	Yes	No	No	No
Vaccination	No	No	No	No	Yes	Yes	Yes
Child has received at least one vaccination							
Outdoor cooking	-0.083*** (0.01)	-0.009*** (0.00)	-0.126*** (0.02)	-0.016*** (0.00)	-0.031** (0.01)	-0.004** (0.00)	-0.007** (0.00)
Observations	122,380	122,380	50,313	50,313	111,945	111,945	38,012
Country dummies	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Year of data collection dummies	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Interview in rainy season dummy	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Household characteristics	No	No	Yes	Yes	No	Yes	Yes
Regular smoking in house	Yes	Yes	Yes	Yes	No	No	No
Vaccination	No	No	No	No	Yes	Yes	Yes

Note: *, **, *** indicate p-values of a 10 percent level, 5 percent level and 1 percent level, respectively. All estimations are clustered on the household level and standard errors are in brackets.

Source: DHS all country dataset from 2005–2014.